



**PCT**

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1

- Claude Monet, F-69110 Ste Foy les LYON (FR). DES-BORDES, Philippe [FR/FR]; 30, Rue Bancel, F-69004 Lyon (FR).

- (74) Common Representative: BAYER CROPSCIENCE AG; Law and Patents, Patents and Licensing, 51368 Leverkusen (DE).**

- (25) **Filing Language:** English

- (26) Publication Language: English

- (30) Priority Data:**  
03015733.3      10 July 2003 (10.07.2003)      EP

- (71) Applicant (for all designated States except US): **BAYER CROPSCIENCE AG** [DE/DE]; Alfred-Nobel-Str. 50, 40789 Monheim (DE).

- (72) Inventors; and

- (75) **Investors/Applicants (for US only):** **DUNKEL, Ralf** [DE/DE]; Krischer Str. 22, 40789 Monheim (DE). **ELBE, Hans-Ludwig** [DE/DE]; Dasnöckel 59, 42329 Wuppertal (DE). **HARTMANN, Benoît** [FR/DE]; Färberstr. 1, 40764 Langenfeld (DE). **GREUL, Jörg, Nico** [DE/DE]; Am Sandberg 30 a, 42799 Leichlingen (DE). **WACHENDORFF-NEUMANN, Ulrike** [DE/DE]; Oberer Markeweg 85, 56566 Neuwied (DE). **DAHMEN, Peter** [DE/DE]; Altebrückerstr. 61, 41470 Neuss (DE). **KUCK, Karl-Helzn** [DE/DE]; Pastor-Löh-Str. 30a, 40764 Langenfeld (DE). **MANSFIELD, Darren, James** [GB/FR]; 25, Montee Bonafous, F-69004 Lyon (FR). **COQUERON, Pierre-Yves** [FR/FR]; 36, Rue de la Tête d'Or, F-69007 Lyon (FR). **RIECK, Heiko** [DE/FR]; 9 rue

- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


- (84) Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

- (54) Title: PYRIDINYLANTIIDS



I

- (57) Abstract:** Novel pyridinylanilides of the formula (I) in which R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and A are as defined in the description, a plurality of processes for preparing these substances and their use for controlling unwanted microorganisms, and also novel intermediates and their preparation.


**WO 2005/004606 A2**

Pyridinylanilides

The present invention relates to novel pyridinylanilides, to several processes for their preparation and to their use for controlling unwanted microorganisms.

It is already known that certain pyridinylanilides have fungicidal properties (cf. WO 01/53259 and JP-A 8-92223). Thus, for example, the pyridinylanilides N-[2-(2-chloro-3-pyridinyl)phenyl]-1,4-dimethyl-1H-pyrrole-3-carboxamide and 1,4-dimethyl-N-{2-[2-(trifluoromethyl)-4-pyridinyl]phenyl}-1H-pyrrole-3-carboxamide (WO 01/53259) or 1-methyl-N-(2-pyridin-2-ylphenyl)-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide and 2-chloro-N-(2-pyridin-2-ylphenyl)nicotinamide (JP-A 8-92223) can be used for the control of fungi. The activity of such compounds, however, is not always satisfactory, particularly if they are applied at low dosages. Other pyridinylanilides, e.g. N-[2-(6-bromo-2-pyridinyl)-4-methylphenyl]-2,2-dimethylpropanamide, N-(4-methyl-2-[6-(trifluoromethyl)-2-pyridinyl]phenyl)cyclopropanecarboxamide and N-{4-methyl-2-[6-(trifluoromethyl)-2-pyridinyl]phenyl}benzamide, are known as herbicides and plant growth regulators (cf. WO 95/09846).

This invention is directed to novel pyridinylanilides of the formula (I)



in which

R represents hydrogen, fluorine, chlorine, methyl or trifluoromethyl;

R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> independently of one another each represents hydrogen, halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy

having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;  
 or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbo-  
 nyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkyl-  
 aminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective  
 5 hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the re-  
 spective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

10  $Q^1$  represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9  
 identical or different halogen atoms, or  $C_3$ - $C_6$ -cycloalkyl and

$Q^2$  represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents  $C_1$ - $C_4$ -alkyl  
 or  $C_1$ - $C_4$ -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl,  
 $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino or phenyl; or  
 represents  $C_2$ - $C_4$ -alkenyloxy or  $C_2$ - $C_4$ -alkynyloxy,

15 represents phenyl, phenoxy, phenylthio, benzoyl, benzoyl-ethenyl, cinnamoyl, heterocyclyl or  
 phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3  
 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substi-  
 tuted, identically or differently, in the ring moiety by halogen, in each case straight-chain or  
 branched  $C_1$ - $C_4$ -alkyl and  $C_1$ - $C_4$ -alkoxy;


20 or

$R^2$  and  $R^3$ , if attached to the pyridinyl moiety in ortho position to each other, furthermore together  
 represent  $C_3$ - $C_4$ -alkylene,  $C_3$ - $C_4$ -alkenylene,  $C_2$ - $C_3$ -oxyalkylene or  $C_1$ - $C_2$ -dioxyalkylene, in  
 each case optionally mono- to tetra-substituted, identically or differently, by fluorine,  
 chlorine, oxo, methyl, ethyl, trifluoromethyl;

25

30  $R^4$  represents hydrogen,  $C_1$ - $C_8$ -alkyl,  $C_1$ - $C_6$ -alkylsulfinyl,  $C_1$ - $C_6$ -alkylsulfonyl,  $C_1$ - $C_4$ -alkoxy- $C_1$ -  
 $C_4$ -alkyl,  $C_3$ - $C_8$ -cycloalkyl;  $C_1$ - $C_6$ -halogenoalkyl,  $C_1$ - $C_4$ -halogenoalkylthio,  $C_1$ - $C_4$ -halogeno-  
 alkylsulfinyl,  $C_1$ - $C_4$ -halogenoalkylsulfonyl, halogeno- $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -halo-  
 genocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; formyl-  
 $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -alkyl)carbonyl- $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -alkoxy)carbonyl- $C_1$ - $C_3$ -alkyl; ( $C_1$ - $C_3$ -  
 halogenoalkyl)carbonyl- $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -halogenoalkoxy)carbonyl- $C_1$ - $C_3$ -alkyl having in  
 each case 1 to 7 fluorine-, chlorine- and/or bromine atoms, ( $C_1$ - $C_3$ -alkyl)carbonyl- $C_1$ - $C_3$ -halo-  
 genoalkyl, ( $C_1$ - $C_3$ -alkoxy)carbonyl- $C_1$ - $C_3$ -halogenoalkyl having in each case 1 to 6 fluorine-,  
 chlorine- and/or bromine atoms, ( $C_1$ - $C_3$ -halogenoalkyl)carbonyl- $C_1$ - $C_3$ -halogenoalkyl, ( $C_1$ - $C_3$ -  
 35 halogenoalkoxy)carbonyl- $C_1$ - $C_3$ -halogenoalkyl having in each case 1 to 13 fluorine-,  
 chlorine- and/or bromine atoms;  $-COR^5$ ,  $-CONR^6R^7$  or  $-CH_2NR^8R^9$ ,

- $R^5$  represents hydrogen,  $C_1$ - $C_8$ -alkyl,  $C_1$ - $C_8$ -alkoxy,  $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -cycloalkyl;  $C_1$ - $C_6$ -halogenoalkyl,  $C_1$ - $C_6$ -halogenoalkoxy, halogeno- $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; or  $-COR^{10}$ ,
- $R^6$  and  $R^7$  independently of one another each represent hydrogen,  $C_1$ - $C_8$ -alkyl,  $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -cycloalkyl;  $C_1$ - $C_6$ -halogenoalkyl, halogeno- $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -halogenocycloalkyl having in each case 1 bis 9 fluorine-, chlorine- and/or bromine atoms,
- $R^6$  and  $R^7$  furthermore together with the nitrogen atom to which they are attached, represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and  $NR^{11}$ , and which heterocycle may optionally be mono- to poly-substituted, identically or differently, by halogen or  $C_1$ - $C_4$ -alkyl,
- $R^8$  and  $R^9$  independently of one another each represent hydrogen,  $C_1$ - $C_8$ -alkyl,  $C_3$ - $C_8$ -cycloalkyl;  $C_1$ - $C_6$ -halogenoalkyl,  $C_3$ - $C_8$ -halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms,
- $R^8$  and  $R^9$  furthermore together with the nitrogen atom to which they are attached, represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and  $NR^{11}$ , and which heterocycle may optionally be mono- to poly-substituted, identically or differently, by halogen or  $C_1$ - $C_4$ -alkyl,
- $R^{10}$  represents hydrogen,  $C_1$ - $C_8$ -alkyl,  $C_1$ - $C_8$ -alkoxy,  $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -cycloalkyl;  $C_1$ - $C_6$ -halogenoalkyl,  $C_1$ - $C_6$ -halogenoalkoxy, halogeno- $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms,
- $R^{11}$  represents hydrogen or  $C_1$ - $C_6$ -alkyl,
- A represents a radical of the formula (A1)



(A1), wherein


- $R^{12}$  represents hydrogen, cyano, halogen, nitro,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_3$ - $C_6$ -cycloalkyl,  $C_1$ - $C_4$ -halogenoalkyl,  $C_1$ - $C_4$ -halogenoalkoxy or  $C_1$ - $C_4$ -halogenoalkylthio each having 1 to 5 halogen atoms, aminocarbonyl or aminocarbonyl- $C_1$ - $C_4$ -alkyl and
- $R^{13}$  represents hydrogen, halogen, cyano,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy or  $C_1$ - $C_4$ -alkylthio and
- $R^{14}$  represents hydrogen,  $C_1$ - $C_4$ -alkyl, hydroxy- $C_1$ - $C_4$ -alkyl,  $C_2$ - $C_6$ -alkenyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_1$ - $C_4$ -alkylthio- $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl,



C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl each having 1 to 5 halogen atoms, or phenyl,

or

A represents a radical of the formula (A2)




(A2), wherein

R<sup>15</sup> and R<sup>16</sup> independently of one another each represent hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms and  
R<sup>17</sup> represents halogen, cyano or C<sub>1</sub>-C<sub>4</sub>-alkyl, or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy each having 1 to 5 halogen atoms,

10 or

A represents a radical of the formula (A3)




(A3), wherein

R<sup>18</sup> and R<sup>19</sup> independently of one another each represent hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms and  
R<sup>20</sup> represents hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms,

15

or

A represents a radical of the formula (A4)




(A4), wherein

R<sup>21</sup> represents hydrogen, halogen, hydroxyl, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy or C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio each having 1 to 5 halogen atoms,

20

or

A represents a radical of the formula (A5)



(A5), wherein

R<sup>22</sup> represents halogen, hydroxyl, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio or C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy each having 1 to 5 halogen atoms and

25

$R^{23}$  represents hydrogen, halogen, cyano,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -halogenoalkyl,  $C_1$ - $C_4$ -halogenoalkoxy each having 1 to 5 halogen atoms,  $C_1$ - $C_4$ -alkylsulphinyl or  $C_1$ - $C_4$ -alkylsulphonyl,

or

5 A represents a radical of the formula (A6)



$R^{24}$  represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms and

$R^{25}$  represents  $C_1$ - $C_4$ -alkyl,

$Q^3$  represents a sulphur or oxygen atom, represents SO, SO<sub>2</sub> or CH<sub>2</sub>,

10 p represents 0, 1 or 2, where  $R^{25}$  represents identical or different radicals if p represents 2,

or

A represents a radical of the formula (A7)



$R^{26}$  represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms,

15 or


A represents a radical of the formula (A8)



$R^{27}$  represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms,

or

20 A represents a radical of the formula (A9)




$R^{28}$  and  $R^{29}$  independently of one another each represent hydrogen, halogen, amino,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms and

$R^{30}$  represents hydrogen, halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms,

25

or

A represents a radical of the formula (A10)



(A10), wherein


R<sup>31</sup> and R<sup>32</sup> independently of one another each represent hydrogen, halogen, amino, nitro, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms and

R<sup>33</sup> represents hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms,

5

or

A represents a radical of the formula (A11)



(A11), wherein


R<sup>34</sup> represents hydrogen, halogen, amino, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di-(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms and

10

R<sup>35</sup> represents halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A12)



(A12), wherein


R<sup>36</sup> represents hydrogen, halogen, amino, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di-(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms and

15

R<sup>37</sup> represents halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A13)




(A13), wherein

R<sup>38</sup> represents halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms,

20

or

A represents a radical of the formula (A14)



(A14), wherein


R<sup>39</sup> represents hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl and

25

$R^{40}$  represents halogen or  $C_1$ - $C_4$ -alkyl,

or

A represents a radical of the formula (A15)




(A15), wherein

5  $R^{41}$  represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A16)




(A16), wherein

10  $R^{42}$  represents hydrogen, halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A17)



(A17), wherein

15  $R^{43}$  represents halogen, hydroxyl,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -halogenoalkyl,  $C_1$ - $C_4$ -halogenoalkylthio or  $C_1$ - $C_4$ -halogenoalkoxy each having 1 to 5 halogen atoms,

excluded compounds of the formula (I), in which


R represents hydrogen and

20  $R^1$ ,  $R^2$  and  $R^3$  independently of one another each represents hydrogen, halogen; or straight-chain or branched alkyl having 1 to 4 carbon atoms; or straight-chain or branched halogenoalkyl having 1 to 4 carbon atoms; and

$R^4$  represents hydrogen

and

25 A represents a radical of the formula (A1)



(A1), wherein


$R^{12}$  represents halogen,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl and

$R^{13}$  represents hydrogen and

$R^{14}$  represents methyl,

or

A represents a radical of the formula (A2)




(A2), wherein

$R^{15}$  and  $R^{16}$  independently of one another each represent hydrogen or  $C_1$ - $C_4$ -alkyl and

$R^{17}$  represents halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl,

or

A represents a radical of the formula (A4)




(A4), wherein

$R^{21}$  represents halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl,

or

A represents a radical of the formula (A5)




(A5), wherein

$R^{22}$  represents halogen and

$R^{23}$  represents hydrogen,

or

A represents a radical of the formula (A6)



(A6), wherein


$R^{24}$  represents methyl and

$Q^3$  represents a sulphur or  $CH_2$ ,

p represents 0,

or

A represents a radical of the formula (A9)




(A9), wherein

$R^{28}$  and  $R^{29}$  independently of one another each represent hydrogen or  $C_1$ - $C_4$ -alkyl and

$R^{30}$  represents methyl,

or

A represents a radical of the formula (A11)




(A11), wherein

5  $R^{34}$  represents hydrogen or  $C_1$ - $C_4$ -alkyl and

$R^{35}$  represents halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl,

or

A represents a radical of the formula (A16)



(A16), wherein

10  $R^{42}$  represents halogen.

The excluded compounds are known from JP-A 8-92223.

In each case the following single compounds known from JP-A 8-92223 are explicitly excluded from the scope of protection:

- 15 2-methyl-N-(2-pyridin-2-ylphenyl)-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide;
- N-(2-pyridin-2-ylphenyl)-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide;
- 4-iodo-N-(2-pyridin-2-ylphenyl)-1,3-thiazole-5-carboxamide;
- 1-methyl-N-(2-pyridin-2-ylphenyl)-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide;
- 3-iodo-1-methyl-N-(2-pyridin-2-ylphenyl)-1H-pyrazole-4-carboxamide;
- 20 2-methyl-N-(2-pyridin-2-ylphenyl)-3-furamide;
- 3-methyl-N-(2-pyridin-2-ylphenyl)thiophene-2-carboxamide;
- 2-chloro-N-(2-pyridin-2-ylphenyl)benzamide;
- 2-chloro-N-(2-pyridin-2-ylphenyl)nicotinamide;
- 3-chloro-N-(2-pyridin-2-ylphenyl)pyrazine-2-carboxamide;
- 25 2-methyl-N-(2-pyridin-2-ylphenyl)-5,6-dihydro-1,4-oxathiine-3-carboxamide;
- 2-chloro-N-(2-pyridin-3-ylphenyl)nicotinamide;
- 2-chloro-N-(2-pyridin-4-ylphenyl)nicotinamide;
- 2-methyl-N-[2-(6-methylpyridin-2-yl)phenyl]-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide;
- 1-methyl-N-[2-(6-methylpyridin-2-yl)phenyl]-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide.

30

The compounds according to the invention may exist in different isomeric forms, in particular in form of stereoisomers, such as for example E- and Z-, threo- and erythro-, optical isomers and optionally in

form of tautomers. The invention relates to all the use of the pure isomers as well as the E- and Z-isomers, the threo- and erythro-isomers, the optical isomers, optional mixtures of these isomers and the possible tautomeric forms.

5 Furthermore, it has been found that pyridinylanilides of the formula (I) are obtained when

a) carboxylic acid derivatives of the formula (II)




in which

$\text{X}^1$  represents halogen or hydroxyl and

10 A is as defined above,

are reacted with amines of the formula (III)




in which  $\text{R}$ ,  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^4$  are as defined above,

15 if appropriate in the presence of a catalyst, if appropriate in the presence of a condensing agent, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

b) halogeno-carboxamides of the formula (IV)




20

in which

$\text{R}$ ,  $\text{R}^4$  and  $\text{A}$  are as defined above, and

$\text{X}^2$  represents bromine or iodine,

25 are reacted with boronic acid derivatives of the formula (V)



in which


$R^1$ ,  $R^2$  and  $R^3$  are as defined above, and

$A^1$  and  $A^2$  each represent hydrogen or together represent tetramethylethylene,

5 in the presence of a catalyst, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

c) carboxamide boronic acid derivatives of the formula (VI)




10 in which

$R$ ,  $R^4$  and  $A$  are as defined above, and

$A^3$  and  $A^4$  each represent hydrogen or together represent tetramethylethylene,

are reacted with pyridinyl derivatives of the formula (VII)




15

in which  $R^1$ ,  $R^2$  and  $R^3$  are as defined above,

in the presence of a catalyst, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

20 d) halogeno-carboxamides of the formula (IV)




in which

$R$ ,  $R^4$  and  $A$  are as defined above, and

$X^2$  represents bromine or iodine,



are reacted with pyridinyl derivatives of the formula (VII)




in which R¹, R² and R³ are as defined above,

in the presence of a palladium or platinum catalyst and in the presence of 4,4',4',5,5,5',5'-octamethyl-2,2'-bis-1,3,2-dioxaborolane [bis(pinacolato)diboron], if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

e) pyridinylanilides of the formula (I-1)



in which R, R¹, R², R³ and A are as defined above,

are reacted with halogenides of the formula (VIII)



in which

X³ represents chlorine, bromine or iodine,

R⁴ᵃ represents C₁-C₈-alkyl, C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₈-cycloalkyl; C₁-C₆-halogenoalkyl, C₁-C₄-halogenoalkylthio, C₁-C₄-halogenoalkylsulfinyl, C₁-C₄-halogenoalkylsulfonyl, halogeno-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₈-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; formyl-C₁-C₃-alkyl, (C₁-C₃-alkyl)carbonyl-C₁-C₃-alkyl, (C₁-C₃-alkoxy)carbonyl-C₁-C₃-alkyl; (C₁-C₃-halogenoalkyl)carbonyl-C₁-C₃-alkyl, (C₁-C₃-halogenoalkoxy)carbonyl-C₁-C₃-alkyl having in each case 1 to 7 fluorine-, chlorine- and/or bromine atoms, (C₁-C₃-alkyl)carbonyl-C₁-C₃-halogenoalkyl, (C₁-C₃-alkoxy)carbonyl-C₁-C₃-halogenoalkyl having in each case 1 to 6 fluorine-, chlorine- and/or bromine atoms, (C₁-C₃-halogenoalkyl)carbonyl-C₁-C₃-halogenoalkyl, (C₁-C₃-halogenoalkoxy)carbonyl-C₁-C₃-halogenoalkyl having in each case 1 to 13 fluorine-, chlorine- and/or bromine atoms; -COR⁵, -CONR⁶R⁷ or -CH₂NR⁸R⁹,

R⁵, R⁶, R⁷, R⁸ and R⁹ are as defined above,

in the presence of a base and in the presence of a diluent.

Finally, it has been found that the novel pyridinylanilides of the formula (I) have very good microbicidal properties and can be used for controlling unwanted microorganisms both in crop protection and in the protection of materials.

- 5 Surprisingly, the pyridinylanilides of the formula (I) according to the invention have considerably better fungicidal activity than the constitutionally most similar active compounds of the prior art having the same direction of action.

The formula (I) provides a general definition of the pyridinylanilides according to the invention.  
10 Preferred definitions of the radicals of the above and/or below mentioned formulae are given in the following. These definitions apply in the same way to the final products of the formula (I) as well as to all intermediates.

- R preferably represents hydrogen.
- 15 R furthermore preferably represents fluorine, which fluorine particularly preferably is placed in 4-, 5- or 6-position, very particularly preferably in 4- or 6-position of the anilide moiety [cf. formula (I) above].
- R furthermore preferably represents chlorine, which chlorine particularly preferably is placed in 5-position of the anilide moiety [cf. formula (I) above].
- 20 R furthermore preferably represents methyl, which methyl particularly preferably is placed in 3-position of the anilide moiety [cf. formula (I) above].
- R furthermore preferably represents trifluoromethyl, which trifluoromethyl particularly preferably is placed in 4- or 5-position of the anilide moiety [cf. formula (I) above].
- 25 R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> independently of one another each preferably represents hydrogen, halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;  
or preferably represents in each case straight-chain or branched alkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 6 carbon atoms;  
or preferably represents in each case straight-chain or branched halogenoalkyl, halogeno-  
30 alkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 4 carbon atoms and 1 to 9 identical or different halogen atoms;  
or preferably represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 4 carbon atoms in the  
35 respective hydrocarbon chain;  
or preferably represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

or preferably represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  preferably represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms or  $C_3$ - $C_6$ -cycloalkyl and

$Q^2$  preferably represents hydroxyl,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -halogenoalkyl or  $C_1$ - $C_4$ -halogenoalkoxy each having 1 to 9 identical or different halogen atoms.

$R^2$  and  $R^3$ , if attached to the pyridinyl moiety in ortho position to each other, furthermore together preferably represent  $-(CH_2)_3-$ ,  $-(CH_2)_4-$ ,  $-CH=CH-CH=CH-$ ,  $-O(CH_2)_2-$ ,  $-O(CH_2)_3-$ ,  $-OCH_2O-$ ,  $-O(CH_2)_2O-$ , in each case optionally mono- to tetra-substituted, identically or differently, by fluorine, chlorine, oxo, methyl, ethyl, trifluoromethyl.

$R^1$ ,  $R^2$  and  $R^3$  independently of one another each particularly preferably represents hydrogen, fluorine, chlorine, bromine, cyano; methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxy, ethoxy, n- or iso-propoxy, n-, iso-, sec- or tert-butoxy, methylthio, ethylthio, n- or iso-propylthio, n-, iso-, sec- or tert-butylthio, trifluoromethyl, trifluoroethyl, difluoromethoxy, trifluoromethoxy, difluorochloromethoxy, trifluoroethoxy, cyclopropyl, cyclopentyl, cyclohexyl, or particularly preferably represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  particularly preferably represents hydrogen, methyl, ethyl, trifluoromethyl or cyclopropyl, and

$Q^2$  particularly preferably represents hydroxyl, methoxy, ethoxy, n-propoxy or isopropoxy.

$R^2$  and  $R^3$ , if attached to the pyridinyl moiety in ortho position to each other, furthermore together particularly preferably represent  $-(CH_2)_3-$ ,  $-(CH_2)_4-$ ,  $-CH=CH-CH=CH-$ ,  $-OCH_2O-$ ,  $-O(CH_2)_2O-$ ,  $-OCF_2O-$ ,  $-O(CF_2)_2O-$ .

$R^1$ ,  $R^2$  and  $R^3$  independently of one another each very particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxy, ethoxy, n- or iso-propoxy, trifluoromethyl, trifluoroethyl, difluoromethoxy, trifluoromethoxy, difluorochloromethoxy, trifluoroethoxy,

or very particularly preferably represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  very particularly preferably represents hydrogen, methyl or ethyl and

$Q^2$  very particularly preferably represents hydroxyl, methoxy, ethoxy, n-propoxy or isopropoxy.

$R^2$  and  $R^3$ , if attached to the pyridinyl moiety in ortho position to each other, furthermore together very particularly preferably represent  $-CH=CH-CH=CH-$ ,  $-OCF_2O-$ ,  $-O(CF_2)_2O-$ .

- R<sup>4</sup> preferably represents hydrogen; C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylsulfonyl, halogeno-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; (C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 7 fluorine-, chlorine- and/or bromine atoms, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl having in each case 1 to 6 fluorine-, chlorine- and/or bromine atoms, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl having in each case 1 to 13 fluorine-, chlorine- and/or bromine atoms; -COR<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>.
- R<sup>4</sup> particularly preferably represents hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, pentyl or hexyl, methylsulfinyl, ethylsulfinyl, n- or iso-propylsulfinyl, n-, iso-, sec- or tert-butylsulfinyl, methylsulfonyl, ethylsulfonyl, n- or iso-propylsulfonyl, n-, iso-, sec- or tert-butylsulfonyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl, trifluoromethyl, trichloromethyl, trifluoroethyl, difluoromethylthio, difluorochloromethylthio, trifluoromethylthio, trifluoromethylsulfinyl, trifluoromethylsulfonyl, trifluoromethoxymethyl; -CH<sub>2</sub>-CHO, -CH<sub>2</sub>CH<sub>2</sub>-CHO, -CH<sub>2</sub>-CO-CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>-CO-CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>-CO-CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>-CO-CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-C(O)OCH<sub>3</sub>, -CH<sub>2</sub>-C(O)OCH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>-C(O)OCH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>-C(O)OCH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>-C(O)OCH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>-C(O)OCH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-CO-CF<sub>3</sub>, -CH<sub>2</sub>-CO-CCl<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CCl<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>-CO-CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>-CO-CH<sub>2</sub>CCl<sub>3</sub>, -CH<sub>2</sub>-C(O)OCH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>-C(O)OCF<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>-C(O)OCH<sub>2</sub>CCl<sub>3</sub>, -CH<sub>2</sub>-C(O)OCCl<sub>2</sub>CCl<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>-C(O)OCH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>-C(O)OCF<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>-C(O)OCH<sub>2</sub>CCl<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>-C(O)O-CCl<sub>2</sub>CCl<sub>3</sub>; -COR<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>.
- R<sup>4</sup> very particularly preferably represents hydrogen; methyl, methoxymethyl, -CH<sub>2</sub>-CHO, -CH<sub>2</sub>CH<sub>2</sub>-CHO, -CH<sub>2</sub>-CO-CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH(CH<sub>3</sub>)<sub>2</sub> or -COR<sup>5</sup>.
- R<sup>5</sup> preferably represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy, halogeno-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; or -COR<sup>10</sup>.
- R<sup>5</sup> particularly preferably represents hydrogen, methyl, ethyl, n- or iso-propyl, tert-butyl, methoxy, ethoxy, tert-butoxy, cyclopropyl; trifluoromethyl, trifluoromethoxy; or -COR<sup>10</sup>.
- R<sup>5</sup> very particularly preferably represents hydrogen, -COCH<sub>3</sub>, -CHO, -COCH<sub>2</sub>OCH<sub>3</sub>, -COCO<sub>2</sub>CH<sub>3</sub>, -COCO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>; or -COR<sup>10</sup>.

- R<sup>6</sup> and R<sup>7</sup> independently of one another each preferably represent hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, halogeno-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms.
- 5 R<sup>6</sup> and R<sup>7</sup> furthermore together with the nitrogen atom to which they are attached, preferably represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and NR<sup>11</sup>, and which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl.
- 10 R<sup>6</sup> and R<sup>7</sup> independently of one another each particularly preferably represent hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl; trifluoromethyl, trichloromethyl, trifluoroethyl, trifluoromethoxymethyl.
- R<sup>6</sup> and R<sup>7</sup> furthermore together with the nitrogen atom to which they are attached, particularly  
15 preferably represent a saturated heterocycle selected from the group consisting of morpholine, thiomorpholine and piperazine, which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by fluorine, chlorine, bromine or methyl and where the piperazine additionally at the second nitrogen atom may be substituted by R<sup>11</sup>.
- 20 R<sup>8</sup> and R<sup>9</sup> independently of one another each preferably represent hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>3</sub>-C<sub>6</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms.
- R<sup>8</sup> and R<sup>9</sup> furthermore together with the nitrogen atom to which they are attached, preferably represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional,  
25 non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and NR<sup>11</sup>, and which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl.
- R<sup>8</sup> and R<sup>9</sup> independently of one another each particularly preferably represent hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl; trifluoromethyl, trichloromethyl,  
30 trifluoroethyl, trifluoromethoxymethyl.
- R<sup>8</sup> and R<sup>9</sup> furthermore together with the nitrogen atom to which they are attached, particularly  
preferably represent a saturated heterocycle selected from the group consisting of morpholine, thiomorpholine and piperazine, which heterocycle may optionally be mono- to  
35 tetra-substituted, identically or differently, by fluorine, chlorine, bromine or methyl and where the piperazine additionally at the second nitrogen atom may be substituted by R<sup>11</sup>.

- R<sup>10</sup> preferably represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy, halogeno-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms.
- 5 R<sup>10</sup> particularly preferably represents hydrogen, methyl, ethyl, n- or iso-propyl, tert-butyl, methoxy, ethoxy, n- or iso-propoxy, tert-butoxy, cyclopropyl; trifluoromethyl, trifluoromethoxy.
- R<sup>11</sup> preferably represents hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl.
- R<sup>11</sup> particularly preferably represents hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl.
- 10 A preferably represents one of the radicals  
A1, A2, A3, A4, A5, A6, A9, A10, A11, A12 or A17.
- A particularly preferably represents one of the radicals  
A1, A2, A4, A5, A6, A9, A11, A17.
- 15 A very particularly preferably represents the radical A1.
- A furthermore very particularly preferably represents the radical A2.
- A furthermore very particularly preferably represents the radical A4.
- A furthermore very particularly preferably represents the radical A5.
- A furthermore very particularly preferably represents the radical A6.
- 20 A furthermore very particularly preferably represents the radical A9.
- A furthermore very particularly preferably represents the radical A11.
- A furthermore very particularly preferably represents the radical A17.
- R<sup>12</sup> preferably represents hydrogen, cyano, fluorine, chlorine, bromine, iodine, methyl, ethyl, iso-propyl, methoxy, ethoxy, methylthio, ethylthio, cyclopropyl, C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>2</sub>-halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms, trifluoromethylthio, difluoromethylthio, aminocarbonyl, aminocarbonylmethyl or aminocarbonylethyl.
- 25 R<sup>12</sup> particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, iso-propyl, monofluoromethyl, monofluoroethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl, dichloromethyl, cyclopropyl, methoxy, ethoxy, trifluoromethoxy, trichloromethoxy, methylthio, ethylthio, trifluoromethylthio or difluoromethylthio and
- 30 R<sup>12</sup> very particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, iso-propyl, monofluoromethyl, monofluoroethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl or trichloromethyl.
- 35

- R<sup>12</sup> especially preferably represents methyl, monofluoromethyl, difluoromethyl or trifluoromethyl.
- 5 R<sup>13</sup> preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, methoxy, ethoxy, methylthio or ethylthio.
- R<sup>13</sup> particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine or methyl.
- R<sup>13</sup> very particularly preferably represents hydrogen, fluorine, chlorine or methyl.
- 10 R<sup>14</sup> preferably represents hydrogen, methyl, ethyl, n-propyl, iso-propyl, C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms, hydroxymethyl, hydroxyethyl, cyclopropyl, cyclopentyl, cyclohexyl or phenyl.
- R<sup>14</sup> particularly preferably represents hydrogen, methyl, ethyl, iso-propyl, trifluoromethyl, difluoromethyl, hydroxymethyl, hydroxyethyl or phenyl.
- R<sup>14</sup> very particularly preferably represents hydrogen, methyl, trifluoromethyl or phenyl.
- 15 R<sup>14</sup> especially preferably represents methyl.
- R<sup>15</sup> and R<sup>16</sup> independently of one another each preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 20 R<sup>15</sup> and R<sup>16</sup> independently of one another each particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl or trichloromethyl.
- R<sup>15</sup> and R<sup>16</sup> independently of one another each very particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, difluoromethyl, trifluoromethyl or trichloromethyl.
- 25 R<sup>15</sup> and R<sup>16</sup> especially preferably each represent hydrogen.
- R<sup>17</sup> preferably represents fluorine, chlorine, bromine, cyano, methyl, ethyl, C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 30 R<sup>17</sup> particularly preferably represents fluorine, chlorine, bromine, cyano, methyl, trifluoromethyl, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy or trichloromethoxy.
- R<sup>17</sup> very particularly preferably represents fluorine, chlorine, bromine, iodine, methyl, trifluoromethyl or trifluoromethoxy.
- R<sup>17</sup> especially preferably represents methyl.

R<sup>18</sup> and R<sup>19</sup> independently of one another each preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

5 R<sup>18</sup> and R<sup>19</sup> independently of one another each particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl or trichloromethyl.

R<sup>18</sup> and R<sup>19</sup> independently of one another each very particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, difluoromethyl, trifluoromethyl or trichloromethyl.

R<sup>18</sup> and R<sup>19</sup> especially preferably each represent hydrogen.

10

R<sup>20</sup> preferably represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

R<sup>20</sup> particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl or trifluoromethyl.

15 R<sup>20</sup> very particularly preferably represents methyl.

R<sup>21</sup> preferably represents hydrogen, fluorine, chlorine, bromine, iodine, hydroxyl, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>2</sub>-halogenoalkoxy or C<sub>1</sub>-C<sub>2</sub>-halogenoalkylthio each having 1 to 5 fluorine, chlorine and/or bromine atoms.

20 R<sup>21</sup> particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, hydroxyl, cyano, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy, trichloromethoxy, trifluoromethylthio, difluoromethylthio, difluorochloromethylthio or trichloromethylthio.

25 R<sup>21</sup> very particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, difluoromethyl, trifluoromethyl or trichloromethyl.

R<sup>21</sup> especially preferably represents iodine, methyl, difluoromethyl or trifluoromethyl.

30 R<sup>22</sup> preferably represents fluorine, chlorine, bromine, iodine, hydroxyl, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio, C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms.

35 R<sup>22</sup> particularly preferably represents fluorine, chlorine, bromine, iodine, hydroxyl, cyano, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl, methoxy, ethoxy, methylthio,



- ethylthio, difluoromethylthio, trifluoromethylthio, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy or trichloromethoxy.
- 5  $R^{22}$  very particularly preferably represents fluorine, chlorine, bromine, iodine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- $R^{23}$  preferably represents hydrogen, fluorine, chlorine, bromine, iodine, cyano,  $C_1$ - $C_4$ -alkyl, methoxy, ethoxy, methylthio, ethylthio,  $C_1$ - $C_2$ -halogenoalkyl or  $C_1$ - $C_2$ -halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms,  $C_1$ - $C_2$ -alkylsulphinyl or  $C_1$ - $C_2$ -alkylsulphonyl.
- 10  $R^{23}$  particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, cyano, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl, methoxy, ethoxy, methylthio, ethylthio, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy, trichloromethoxy, methylsulphinyl or methylsulphonyl.
- 15  $R^{23}$  very particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, trichloromethyl, methylsulphinyl or methylsulphonyl.
- $R^{23}$  especially preferably represents hydrogen.
- 20  $R^{24}$  preferably represents methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- $R^{24}$  particularly preferably represents methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- 25  $R^{25}$  preferably represents methyl or ethyl.
- $R^{25}$  particularly preferably represents methyl.
- $Q^3$  preferably represents a sulphur atom,  $SO_2$  or  $CH_2$ .
- $Q^3$  particularly preferably represents a sulphur atom or  $CH_2$ .
- 30  $Q^3$  very particularly preferably represents a sulphur atom.
- p preferably represents 0 or 1.
- p particularly preferably represents 0.
- 35  $R^{26}$  preferably represents methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

R<sup>26</sup> particularly preferably represents methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

R<sup>26</sup> very particularly preferably represents methyl, trifluoromethyl, difluoromethyl or trichloromethyl.

5

R<sup>27</sup> preferably represents methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

R<sup>27</sup> particularly preferably represents methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

10 R<sup>27</sup> very particularly preferably represents methyl, trifluoromethyl, difluoromethyl or trichloromethyl.

R<sup>28</sup> and R<sup>29</sup> independently of one another each preferably represent hydrogen, fluorine, chlorine, bromine, amino, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

15

R<sup>28</sup> and R<sup>29</sup> independently of one another each particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

R<sup>28</sup> and R<sup>29</sup> independently of one another each very particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.

20

R<sup>28</sup> and R<sup>29</sup> especially preferably each represent hydrogen.

R<sup>30</sup> preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

25

R<sup>30</sup> particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

R<sup>30</sup> very particularly preferably represents hydrogen, fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.

R<sup>30</sup> especially preferably represents methyl.

30

R<sup>31</sup> and R<sup>32</sup> independently of one another each preferably represent hydrogen, fluorine, chlorine, bromine, amino, nitro, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

R<sup>31</sup> and R<sup>32</sup> independently of one another each particularly preferably represent hydrogen, fluorine, chlorine, bromine, nitro, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

35

R<sup>31</sup> and R<sup>32</sup> independently of one another each very particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.

R<sup>31</sup> and R<sup>32</sup> especially preferably each represent hydrogen.

5 R<sup>33</sup> preferably represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

R<sup>33</sup> particularly preferably represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

10 R<sup>33</sup> very particularly preferably represents hydrogen, fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.

R<sup>33</sup> especially preferably represents methyl.

15 R<sup>34</sup> preferably represents hydrogen, fluorine, chlorine, bromine, amino, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino, cyano, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

R<sup>34</sup> particularly preferably represents hydrogen, fluorine, chlorine, bromine, amino, methyl-amino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

20 R<sup>34</sup> very particularly preferably represents hydrogen, fluorine, chlorine, bromine, amino, methylamino, dimethylamino, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.

R<sup>34</sup> especially preferably represents amino, methylamino, dimethylamino, methyl or trifluoromethyl.

25 R<sup>35</sup> preferably represents fluorine, chlorine, bromine, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

R<sup>35</sup> particularly preferably represents fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

R<sup>35</sup> very particularly preferably represents fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.

30 R<sup>35</sup> especially preferably represents methyl, trifluoromethyl or difluoromethyl.

R<sup>36</sup> preferably represents hydrogen, fluorine, chlorine, bromine, amino, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino, cyano, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

- R<sup>36</sup> particularly preferably represents hydrogen, fluorine, chlorine, bromine, amino, methylamino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- 5 R<sup>36</sup> very particularly preferably represents hydrogen, fluorine, chlorine, bromine, amino, methylamino, dimethylamino, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R<sup>36</sup> especially preferably represents amino, methylamino, dimethylamino, methyl or trifluoromethyl.
- 10 R<sup>37</sup> preferably represents fluorine, chlorine, bromine, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- R<sup>37</sup> particularly preferably represents fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- R<sup>37</sup> very particularly preferably represents fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- 15 R<sup>37</sup> especially preferably represents methyl, trifluoromethyl or difluoromethyl.
- R<sup>38</sup> preferably represents fluorine, chlorine, bromine, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 20 R<sup>38</sup> particularly preferably represents fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- R<sup>38</sup> very particularly preferably represents fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R<sup>39</sup> preferably represents hydrogen, methyl or ethyl.
- 25 R<sup>39</sup> particularly preferably represents methyl.
- R<sup>40</sup> preferably represents fluorine, chlorine, bromine, methyl or ethyl,
- R<sup>40</sup> particularly preferably represents fluorine, chlorine or methyl.
- 30 R<sup>41</sup> preferably represents methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- R<sup>41</sup> particularly preferably represents methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- R<sup>41</sup> very particularly preferably represents methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- 35 R<sup>41</sup> especially preferably represents methyl or trifluoromethyl.

$R^{42}$  preferably represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

$R^{42}$  particularly preferably represents hydrogen, fluorine, chlorine, bromine, methyl or trifluoromethyl.

5


$R^{43}$  preferably represents fluorine, chlorine, bromine, iodine, hydroxyl,  $C_1$ - $C_4$ -alkyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio,  $C_1$ - $C_2$ -halogenoalkyl or  $C_1$ - $C_2$ -halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms.

$R^{43}$  particularly preferably represents fluorine, chlorine, bromine, iodine, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl.

10

$R^{43}$  very particularly preferably represents fluorine, chlorine, bromine, iodine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.

15 Moreover, emphasis is given to compounds of the formula (I-1)




in which R,  $R^1$ ,  $R^2$ ,  $R^3$  and A are as defined above,  
excluded compounds of the formula (I-1), in which

R represents hydrogen and

20  $R^1$ ,  $R^2$  and  $R^3$  independently of one another each represents hydrogen, halogen;  
or represents straight-chain or branched alkyl having 1 to 4 carbon atoms;  
or represents straight-chain or branched halogenoalkyl having 1 to 4 carbon atoms;

and

A represents a radical of the formula (A1)



(A1), wherein

25


$R^{12}$  represents halogen,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl and

$R^{13}$  represents hydrogen and

$R^{14}$  represents methyl,

or

A represents a radical of the formula (A2)




(A2), wherein

R<sup>15</sup> and R<sup>16</sup> independently of one another each represent hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl and  
R<sup>17</sup> represents halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl,

5 or

A represents a radical of the formula (A4)




(A4), wherein

R<sup>21</sup> represents halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl,

or

10 A represents a radical of the formula (A5)




(A5), wherein

R<sup>22</sup> represents halogen and

R<sup>23</sup> represents hydrogen,

or

15 A represents a radical of the formula (A6)



(A6), wherein


R<sup>24</sup> represents methyl and

Q<sup>3</sup> represents a sulphur or CH<sub>2</sub>,

p represents 0,

20 or

A represents a radical of the formula (A9)




(A9), wherein

R<sup>28</sup> and R<sup>29</sup> independently of one another each represent hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl and

R<sup>30</sup> represents methyl,

25 or

A represents a radical of the formula (A11)




(A11), wherein

$R^{34}$  represents hydrogen or  $C_1$ - $C_4$ -alkyl and

$R^{35}$  represents halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl,

or


5 A represents a radical of the formula (A16)



(A16), wherein

$R^{42}$  represents halogen.

Moreover, emphasis is given to compounds of the formula (I-2)



(I-2)

10

in which  $R$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^{4a}$  and  $A$  are as defined above.


$R^{4a}$  preferably represents  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_4$ -alkylsulfinyl,  $C_1$ - $C_4$ -alkylsulfonyl,  $C_1$ - $C_3$ -alkoxy- $C_1$ - $C_3$ -alkyl,  $C_3$ - $C_6$ -cycloalkyl;  $C_1$ - $C_4$ -halogenoalkyl,  $C_1$ - $C_4$ -halogenoalkylthio,  $C_1$ - $C_4$ -halogenoalkylsulfinyl,  $C_1$ - $C_4$ -halogenoalkylsulfonyl, halogeno- $C_1$ - $C_3$ -alkoxy- $C_1$ - $C_3$ -alkyl,  $C_3$ - $C_6$ -halogenocycloalkyl each having 1 to 9 fluorine-, chlorine- and/or bromine atoms; formyl- $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -alkyl)carbonyl- $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -alkoxy)carbonyl- $C_1$ - $C_3$ -alkyl; ( $C_1$ - $C_3$ -halogenoalkyl)carbonyl- $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -halogenoalkoxy)carbonyl- $C_1$ - $C_3$ -alkyl having in each case 1 to 7 fluorine-, chlorine- and/or bromine atoms, ( $C_1$ - $C_3$ -alkyl)carbonyl- $C_1$ - $C_3$ -halogenoalkyl, ( $C_1$ - $C_3$ -alkoxy)carbonyl- $C_1$ - $C_3$ -halogenoalkyl having in each case 1 to 6 fluorine-, chlorine- and/or bromine atoms, ( $C_1$ - $C_3$ -halogenoalkyl)carbonyl- $C_1$ - $C_3$ -halogenoalkyl, ( $C_1$ - $C_3$ -halogenoalkoxy)carbonyl- $C_1$ - $C_3$ -halogenoalkyl having in each case 1 to 13 fluorine-, chlorine- and/or bromine atoms;  $-COR^5$ ,  $-CONR^6R^7$  or  $-CH_2NR^8R^9$ , where  $R^5$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined above.

25  $R^{4a}$  particularly preferably represents methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, pentyl or hexyl, methylsulfinyl, ethylsulfinyl, n- or iso-propylsulfinyl, n-, iso-, sec- or tert-butylsulfinyl, methylsulfonyl, ethylsulfonyl, n- or iso-propylsulfonyl, n-, iso-, sec- or tert-butylsulfonyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclo-

pentyl, cyclohexyl, trifluoromethyl, trichloromethyl, trifluoroethyl, difluoromethylthio, difluorochloromethylthio, trifluoromethylthio, trifluoromethylsulfinyl, trifluoromethylsulfonyl, trifluoromethoxymethyl;  $-\text{CH}_2\text{-CHO}$ ,  $-\text{CH}_2\text{CH}_2\text{-CHO}$ ,  $-\text{CH}_2\text{-CO-CH}_3$ ,  $-\text{CH}_2\text{-CO-CH}_2\text{CH}_3$ ,  $-\text{CH}_2\text{-CO-CH(CH}_3)_2$ ,  $-\text{CH}_2\text{CH}_2\text{-CO-CH}_3$ ,  $-\text{CH}_2\text{CH}_2\text{-CO-CH}_2\text{CH}_3$ ,  $-\text{CH}_2\text{CH}_2\text{-CO-CH(CH}_3)_2$ ,  $-\text{CH}_2\text{-C(O)OCH}_3$ ,  $-\text{CH}_2\text{-C(O)OCH}_2\text{CH}_3$ ,  $-\text{CH}_2\text{-C(O)OCH(CH}_3)_2$ ,  $-\text{CH}_2\text{CH}_2\text{-C(O)OCH}_3$ ,  $-\text{CH}_2\text{CH}_2\text{-C(O)OCH}_2\text{CH}_3$ ,  $-\text{CH}_2\text{CH}_2\text{-C(O)OCH(CH}_3)_2$ ,  $-\text{CH}_2\text{-CO-CF}_3$ ,  $-\text{CH}_2\text{-CO-CCl}_3$ ,  $-\text{CH}_2\text{-CO-CH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{-CO-CH}_2\text{CCl}_3$ ,  $-\text{CH}_2\text{CH}_2\text{-CO-CH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CH}_2\text{-CO-CH}_2\text{CCl}_3$ ,  $-\text{CH}_2\text{-C(O)OCH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{-C(O)OCF}_2\text{CF}_3$ ,  $-\text{CH}_2\text{-C(O)OCH}_2\text{CCl}_3$ ,  $-\text{CH}_2\text{-C(O)OCCl}_2\text{CCl}_3$ ,  $-\text{CH}_2\text{CH}_2\text{-C(O)OCH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CH}_2\text{-C(O)OCF}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CH}_2\text{-C(O)OCH}_2\text{CCl}_3$ ,  $-\text{CH}_2\text{CH}_2\text{-C(O)O-CCl}_2\text{CCl}_3$ ;  $-\text{COR}^5$ ,  $-\text{CONR}^6\text{R}^7$  or  $-\text{CH}_2\text{NR}^8\text{R}^9$ , where  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$ ,  $\text{R}^8$  and  $\text{R}^9$  are as defined above.


$\text{R}^{4a}$  very particularly preferably represents methyl, methoxymethyl,  $-\text{CH}_2\text{-CHO}$ ,  $-\text{CH}_2\text{CH}_2\text{-CHO}$ ,  $-\text{CH}_2\text{-CO-CH}_3$ ,  $-\text{CH}_2\text{-CO-CH}_2\text{CH}_3$ ,  $-\text{CH}_2\text{-CO-CH(CH}_3)_2$  or  $-\text{COR}^5$ , where  $\text{R}^5$  is as defined above.

Moreover, emphasis is given to compounds of the formula (I-3)



in which  $\text{R}$ ,  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^4$  and  $\text{A}$  are as defined above,  
where 2-chloro-N-(2-pyridin-3-ylphenyl)nicotinamide is excluded.


Moreover, emphasis is given to compounds of the formula (I-4)



in which  $\text{R}$ ,  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^4$  and  $\text{A}$  are as defined above,  
where 2-chloro-N-(2-pyridin-4-ylphenyl)nicotinamide is excluded.

Moreover, emphasis is given to compounds of the formula (I-5)






in which R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and A are as defined above,

where 2-methyl-N-(2-pyridin-2-ylphenyl)-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide; N-(2-pyridin-2-ylphenyl)-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide; 4-iodo-N-(2-pyridin-2-ylphenyl)-1,3-thiazole-5-carboxamide; 1-methyl-N-(2-pyridin-2-ylphenyl)-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide; 3-iodo-1-methyl-N-(2-pyridin-2-ylphenyl)-1H-pyrazole-4-carboxamide; 2-methyl-N-(2-pyridin-2-ylphenyl)-3-furamide; 3-methyl-N-(2-pyridin-2-ylphenyl)thiophene-2-carboxamide; 2-chloro-N-(2-pyridin-2-ylphenyl)benzamide; 2-chloro-N-(2-pyridin-2-ylphenyl)nicotinamide; 3-chloro-N-(2-pyridin-2-ylphenyl)pyrazine-2-carboxamide; 2-methyl-N-(2-pyridin-2-ylphenyl)-5,6-dihydro-1,4-oxathiine-3-carboxamide; 2-methyl-N-[2-(6-methylpyridin-2-yl)phenyl]-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide; 1-methyl-N-[2-(6-methylpyridin-2-yl)phenyl]-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide are excluded.


Moreover, emphasis is given to compounds of the formula (I-6)



in which R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and A are as defined above,

where 2-chloro-N-(2-pyridin-3-ylphenyl)nicotinamide is excluded.


Moreover, emphasis is given to compounds of the formula (I-7)



in which R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and A are as defined above,

where 2-chloro-N-(2-pyridin-4-ylphenyl)nicotinamide is excluded.


Moreover, emphasis is given to compounds of the formula (I-8)



in which R, R¹, R², R³ and A are as defined above,


where 2-methyl-N-(2-pyridin-2-ylphenyl)-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide; N-(2-pyridin-2-ylphenyl)-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide; 4-iodo-N-(2-pyridin-2-ylphenyl)-1,3-thiazole-5-carboxamide; 1-methyl-N-(2-pyridin-2-ylphenyl)-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide; 3-iodo-1-methyl-N-(2-pyridin-2-ylphenyl)-1H-pyrazole-4-carboxamide; 2-methyl-N-(2-pyridin-2-ylphenyl)-3-furamide; 3-methyl-N-(2-pyridin-2-ylphenyl)thiophene-2-carboxamide; 2-chloro-N-(2-pyridin-2-ylphenyl)benzamide; 2-chloro-N-(2-pyridin-2-ylphenyl)nicotinamide; 3-chloro-N-(2-pyridin-2-ylphenyl)pyrazine-2-carboxamide; 2-methyl-N-(2-pyridin-2-ylphenyl)-5,6-dihydro-1,4-oxathine-3-carboxamide; 2-methyl-N-[2-(6-methylpyridin-2-yl)phenyl]-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide; 1-methyl-N-[2-(6-methylpyridin-2-yl)phenyl]-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide are excluded.

Moreover, emphasis is given to compounds of the formula (I-9)




in which R, R¹, R², R³, R⁴ᵃ and A are as defined above.

Moreover, emphasis is given to compounds of the formula (I-10)




in which R, R¹, R², R³, R⁴ᵃ and A are as defined above.

Moreover, emphasis is given to compounds of the formula (I-11)




in which R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4a</sup> and A are as defined above.

5 Moreover, emphasis is given to compounds of the formula (I-12)



in which R, R<sup>4</sup> and A are as defined above, and R<sup>1a</sup> is as defined below.


Moreover, emphasis is given to compounds of the formula (I-13)



10

in which R, R<sup>4</sup> and A are as defined above, and R<sup>1a</sup> is as defined below.


Moreover, emphasis is given to compounds of the formula (I-14)



15


in which R, R<sup>4</sup> and A are as defined above, and R<sup>1a</sup> and R<sup>2a</sup> are as defined below.

Moreover, emphasis is given to compounds of the formula (I-15)



in which R, R<sup>4</sup> and A are as defined above, and R<sup>1a</sup> and R<sup>2a</sup> are as defined below.

5 Moreover, emphasis is given to compounds of the formula (I-16)



in which R, R<sup>4</sup> and A are as defined above, and R<sup>1a</sup>, R<sup>2a</sup> and R<sup>3a</sup> are as defined below.

- R<sup>1a</sup>, R<sup>2a</sup> and R<sup>3a</sup> independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;
- 10 or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;
- or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each
- 15 case 2 to 6 carbon atoms;
- or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;
- or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy
- 20 having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;
- or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxy carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the
- 25 respective hydrocarbon chain;
- or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms, or  $C_3$ - $C_6$ -cycloalkyl and

$Q^2$  represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino or phenyl; or represents  $C_2$ - $C_4$ -alkenyloxy or  $C_2$ - $C_4$ -alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoyl-ethenyl, cinnamoyl, heterocycl-yl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocycl-ylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched  $C_1$ - $C_4$ -alkyl and  $C_1$ - $C_4$ -alkoxy.

$R^{1a}$ ,  $R^{2a}$  and  $R^{3a}$  independently of one another each preferably represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl; or preferably represents in each case straight-chain or branched alkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 6 carbon atoms;

or preferably represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 4 carbon atoms and 1 to 9 identical or different halogen atoms;

or preferably represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 4 carbon atoms in the respective hydrocarbon chain;

or preferably represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms; or preferably represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  preferably represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms or  $C_3$ - $C_6$ -cycloalkyl and

$Q^2$  preferably represents hydroxyl,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -halogenoalkyl or  $C_1$ - $C_4$ -halogenoalkoxy each having 1 to 9 identical or different halogen atoms.

$R^{1a}$ ,  $R^{2a}$  and  $R^{3a}$  independently of one another each particularly preferably represents fluorine, chlorine, bromine, cyano; methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxy, ethoxy, n- or iso-propoxy, n-, iso-, sec- or tert-butoxy, methylthio, ethylthio, n- or iso-propylthio, n-, iso-, sec- or tert-butylthio, trifluoromethyl, trifluoroethyl, difluoromethoxy, trifluor-

omethoxy, difluorochloromethoxy, trifluoroethoxy, cyclopropyl, cyclopentyl, cyclohexyl, or particularly preferably represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  particularly preferably represents hydrogen, methyl, ethyl, trifluoromethyl or cyclopropyl, and

5  $Q^2$  particularly preferably represents hydroxyl, methoxy, ethoxy, n-propoxy or isopropoxy.

$R^{1a}$ ,  $R^{2a}$  and  $R^{3a}$  independently of one another each very particularly preferably represents fluorine, chlorine, bromine, iodine, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxy, 10 ethoxy, n- or iso-propoxy, trifluoromethyl, trifluoroethyl, difluoromethoxy, trifluoromethoxy, difluorochloromethoxy, trifluoroethoxy,

or very particularly preferably represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  very particularly preferably represents hydrogen, methyl or ethyl and

15  $Q^2$  very particularly preferably represents hydroxyl, methoxy, ethoxy, n-propoxy or isopropoxy.

Saturated or unsaturated hydrocarbon radicals, such as alkyl or alkenyl, can in each case be straight-chain or branched as far as this is possible, even in combination with heteroatoms, such as, for example, in alkoxy.

20 Optionally substituted radicals can be mono- or poly-substituted, where in the case of poly-substitutions the substituents can be identical or different.

Halogen-substituted radicals, such as, for example, halogenoalkyl, are mono- or poly-halogenated. In the case of poly-halogenation, the halogen atoms can be identical or different. Here, halogen represents fluorine, chlorine, bromine and iodine, in particular fluorine, chlorine and bromine.


25 However, the general or preferred radical definitions or illustrations listed above can also be combined with one another as desired, i.e. including combinations between the respective ranges and preferred ranges. They apply to the end products and, correspondingly, to precursors and intermediates. Moreover, individual definitions may not apply.

30

#### Detailed Description of the Processes and Intermediates

##### **Process (a)**

35 Using 2-(trifluoromethyl)benzoyl chloride and 2-[3-fluoro-5-(trifluoromethyl)-2-pyridinyl]phenylamine as starting materials, the course of the process (a) according to the invention can be illustrated by the formula scheme below.



The formula (II) provides a general definition of the carboxylic acid derivatives required as starting materials for carrying out the process (a) according to the invention. In this formula, A preferably has those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred or very particularly preferred for these radicals.  $\text{X}^1$  preferably represents chlorine, bromine or hydroxyl, particularly preferably chlorine or hydroxyl.


The carboxylic acid derivatives of the formula (II) are known or can be prepared by known processes (cf. WO 93/11117, EP-A 0 545 099, EP-A 0 589 301 and EP-A 0 589 313).

The formula (III) provides a general definition of the amines required as reaction components for carrying out the process (a) according to the invention. In this formula, R,  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  preferably have those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred or very particularly preferred for these radicals.

The amines of the formula (III) excluded compounds of the formula (III), in which R represents hydrogen and  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  independently of one another each represents hydrogen, halogen, straight-chain or branched alkyl having 1 to 4 carbon atoms or straight-chain or branched halogenoalkyl having 1 to 4 carbon atoms; and  $\text{R}^4$  represents hydrogen are novel (cf. JP-A 8-92223). Amines of the formula (III) employed for the production of compounds of the formulae (I-12), (I-13), (I-14), (I-15) and (I-16) are novel as well. Some of them can be prepared by known methods (Heterocycles 1989, 29, 1013-1016; J. Med. Chem. 1996, 39, 892-903; Synthesis 1995, 713-16; Synth. Commun. 1994, 24, 267-272; DE-A 27 27 416; Synthesis 1994, 142-144; EP-A 0 824 099; WO 93/11117, EP-A 0 545 099, EP-A 0 589 301, EP-A 0 589 313 and WO 02/38542).

Moreover, aniline derivatives of the formula (III) are obtained by

f) reacting 2-halo-amines of the general formula (IX)




in which

R and R<sup>4</sup> are as defined above and

Hal represents halogen,

5

with boronic acid derivatives of the formula (V)




in which R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, A<sup>1</sup> and A<sup>2</sup> are as defined above,

if appropriate in the presence of an acid binder, and if appropriate in the presence of an inert organic diluent, and if appropriate in the presence of a catalyst,

10

or

g) reacting boronic acid derivatives of the formula (X)




in which

15

R and R<sup>4</sup> are as defined above, and

A<sup>5</sup> and A<sup>6</sup> each represent hydrogen or together represent tetramethylethylene,

with pyridinyl derivatives of the formula (VII)



20


in which R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are as defined above,

if appropriate in the presence of an acid binder, and if appropriate in the presence of an inert organic diluent, and if appropriate in the presence of a catalyst,

or

h) reacting 2-halo-amines of the general formula (IX)






in which

R and R<sup>4</sup> are as defined above and

Hal represents halogen,

5

with pyridinyl derivatives of the formula (VII)



in which R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are as defined above,

in the presence of a palladium or platinum catalyst and in the presence of 4,4,4',4',5,5,5',5'-  
 10 octamethyl-2,2'-bis-1,3,2-dioxaborolane, if appropriate in the presence of an acid binder and  
 if appropriate in the presence of a diluent.

The formula (IX) provides a general definition of the 2-halo-amines required as reaction components  
 for carrying out the processes (f) and (h) according to the invention. In this formula R and R<sup>4</sup>  
 15 preferably have those meanings which have already been mentioned in connection with the  
 description of the compounds of the formula (I) according to the invention as being preferred,  
 particularly preferred or very particularly preferred for these radicals. Hal preferably represents  
 chlorine, bromine or iodine, particularly preferably bromine or iodine.

20 2-Halo-amines of the formula (IX) are known and/or can be prepared by known methods from the  
 corresponding nitro compounds by reduction. In the case, that R<sup>4</sup> does not represent hydrogen, the  
 compounds of formula (IX) can be obtained by known derivatizations of the resulting aniline  
 derivatives.

The boronic acid derivatives of the formula (V) furthermore required as starting materials for  
 25 carrying out the process (f) according to the invention are described in more detail below in  
 connection with the process (b) according to the invention.

The formula (X) provides a general definition of the boronic acid derivatives required as reaction  
 components for carrying out the process (g) according to the invention. In this formula R and R<sup>4</sup>  
 30 preferably have those meanings which have already been mentioned in connection with the  
 description of the compounds of the formula (I) according to the invention as being preferred,


particularly preferred or very particularly preferred for these radicals. A<sup>5</sup> and A<sup>6</sup> preferably each represent hydrogen or together represent tetramethylethylene.

The boronic acid derivatives of the formula (X) are known and/or can be obtained by known methods.

The phenyl derivatives of the formula (VII) furthermore required as starting materials for carrying out the processes (g) and (h) according to the invention are illustrated in more detail below, in connection with the process (c) according to the invention.

#### Process (b)

Using *N*-(2-bromophenyl)-3-(difluoromethyl)-1-methyl-1*H*-pyrazole-4-carboxamide and 2-chloro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine as starting materials and a catalyst, the course of the process (b) according to the invention can be illustrated by the formula scheme below.



The formula (IV) provides a general definition of the halogeno-carboxamides required as starting materials for carrying out the process (b) according to the invention. In this formula R, R<sup>4</sup> and A preferably have those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred or very particularly preferred for these radicals.

The carboxamide derivatives of the formula (IV) are known or can be prepared by known processes (cf. WO 91/01311, EP-A 0 371 950). They are obtained, for example, by

- i) reacting carboxylic acid derivatives of the formula (II)




in which

X<sup>1</sup> represents halogen or hydroxyl and

A is as defined above,

with 2-halo-amines of the general formula (IX)



in which

R and R<sup>4</sup> are as defined above and

- 5      Hal      represents halogen,  
 if appropriate in the presence of a catalyst, if appropriate in the presence of a condensing agent, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent.


- 10    The carboxylic acid derivatives of the formula (III) required as starting materials for carrying out the process (i) according to the invention are illustrated in more detail above, in connection with the process (a) according to the invention.

- 15    The 2-halo-amines of the formula (IX) furthermore required as starting materials for carrying out the process (i) according to the invention are illustrated in more detail above, in connection with the process (f) according to the invention.

- 20    The formula (V) provides a general definition of the boronic acid derivatives furthermore required as starting materials for carrying out the process (b) according to the invention. In this formula R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> preferably have those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred or very particularly preferred for these radicals. A<sup>1</sup> and A<sup>2</sup> preferably each represent hydrogen or together represent tetramethylethylene.

- 25    The boronic acid derivatives of the formula (V) are known and/or can be prepared by known processes (cf. WO 01/90084 and US 5,633,218). They are obtained, for example, by

- k)      reacting pyridinyl derivatives of the formula (VII)



in which R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are as defined above,

30

with boric acid esters of the formula (XI)



in which Alk represents C<sub>1</sub>-C<sub>4</sub>-alkyl,

- 5 or with 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bis-1,3,2-dioxaborolane  
in the presence of magnesium or alkyllithium, if appropriate in the presence of a diluent (for  
example tetrahydrofuran).


The formula (XI) provides a general definition of the boric acid esters required as reaction  
10 components for carrying out the process (h) according to the invention. In this formula, Alk  
preferably represents methyl, ethyl, n- or iso-propyl, particularly preferably methyl or ethyl.

The boric acid esters of the formula (XI) are known chemicals for synthesis.

- 15 The pyridinyl derivatives of the formula (VII) furthermore required as starting materials for carrying  
out the process (h) according to the invention are illustrated in more detail below, in connection with  
the process (c) according to the invention.

#### Process (c)

- 20 Using 2-[[[(3-methyl-2-thienyl)carbonyl]amino]phenyl]boronic acid and 2-bromo-3-chloro-5-(  
(trifluoromethyl)pyridine as starting materials and a catalyst, the course of the process (c) according  
to the invention can be illustrated by the formula scheme below.




- The formula (VI) provides a general definition of the carboxamide boronic acid derivatives required  
25 as reaction components for carrying out the process (c) according to the invention. In this formula R,  
R<sup>4</sup> and A preferably have those meanings which have already been mentioned in connection with the  
description of the compounds of the formula (I) according to the invention as being preferred, parti-  
cularly preferred or very particularly preferred for these radicals. A<sup>3</sup> and A<sup>4</sup> preferably each represent  
hydrogen or together represent tetramethylethylene.

The carboxamide boronic acid derivatives of the formula (VI) are known and/or can be prepared by known processes.

- 5 The formula (VII) provides a general definition of the pyridinyl derivatives required as starting materials for carrying out the process (c) according to the invention. In this formula  $R^1$ ,  $R^2$  and  $R^3$  preferably have those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred or very particularly preferred for these radicals.
- 10 The pyridinyl derivatives of the formula (VII) are known or can be prepared by known processes (cf. Synth. Commun. 2000, 30, 665-669, Synth. Commun. 1999, 29, 1697-1701, and cf. also the examples below).

#### Process (d)

- 15 Using *N*-(2-bromophenyl)-2-chloronicotinamide and 2-bromo-5-chloropyridine as starting materials and a catalyst and 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bis-1,3,2-dioxaborolane, the course of the process (d) according to the invention can be illustrated by the formula scheme below.




- 20 The halogeno-carboxamides of the formula (IV) and the pyridinyl derivatives of the formula (VII) required as starting materials for carrying out the process (d) according to the invention are already described above in connection with the processes (b) and (c) according to the invention.

4,4,4',4',5,5,5',5'-Octamethyl-2,2'-bis-1,3,2-dioxaborolane furthermore required for carrying out process (d) according to the invention is a known chemical substance.

25

#### Process (e)

Using *N*-[2-(5-chloro-2-pyridinyl)phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide and acetyl chloride as starting materials, the course of the process (e) according to the invention can be illustrated by the formula scheme below.



The formula (I-1) provides a general definition of the pyridinylanilides required as starting materials for carrying out the process (e) according to the invention. In this formula R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and A preferably have those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred or very particularly preferred for these radicals.

The compounds of the formula (I-1) are compounds according to the invention and can be obtained according to any of the processes (a) to (d).

The formula (VIII) provides a general definition of the halogenides required as starting materials for carrying out the process (e) according to the invention. In this formula R<sup>4a</sup> preferably has those meanings which have already been mentioned in connection with the description of the compounds of the formula (I-2) according to the invention as being preferred, particularly preferred or very particularly preferred for these radicals. X<sup>3</sup> represents chlorine, bromine or iodine.

Halogenides of the formula (VIII) are widely known.

#### Reaction Conditions

Suitable diluents for carrying out the processes (a) and (i) according to the invention are all customary inert organic solvents. Preference is given to using aliphatic, alicyclic or aromatic hydrocarbons, such as petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin; halogenated hydrocarbons, such as chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichloroethane or trichloroethane; ethers, such as diethyl ether, diisopropyl ether, methyl tert-butyl ether, methyl tert-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole; nitriles, such as acetonitrile, propionitrile, n- or iso-butyronitrile or benzonitrile; amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide; mixtures thereof with water or pure water.

Suitable diluents for carrying out the processes (b), (c), (d), (f), (g) and (h) according to the invention are in each case all customary inert organic solvents. Preference is given to using aliphatic, alicyclic

or aromatic hydrocarbons, such as petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin; ethers, such as diethyl ether, diisopropyl ether, methyl tert-butyl ether, methyl tert-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole; nitriles, such as acetonitrile, propionitrile, n- or iso-butyronitrile or benzonitrile; amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide; esters, such as methyl acetate or ethyl acetate; sulphoxides, such as dimethylsulphoxide; or sulphones, such as sulfolane; alcohols, such as methanol, ethanol, n- or iso-propanol, n-, iso-, sec- or tert-butanol, ethanediol, propane-1,2-diol, ethoxyethanol, methoxyethanol, diethyleneglycolmonomethylether, diethyleneglycolmonoethylether; mixtures thereof with water or pure water.

Suitable diluents for carrying out the process (e) according to the invention are all customary inert organic solvents. Preference is given to using aliphatic, alicyclic or aromatic hydrocarbons, such as petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin; halogenated hydrocarbons, such as chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichloroethane or trichloroethane; ethers, such as diethyl ether, diisopropyl ether, methyl tert-butyl ether, methyl tert-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole; amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide.

Suitable acid binders for carrying out the processes (a) and (i) according to the invention are all inorganic and organic bases customary for such reactions. Preference is given to using alkaline earth metal or alkali metal hydrides, hydroxides, amides, alcoholates, acetates, carbonates or hydrogen carbonates, such as sodium hydride, sodium amide, lithium diisopropylamide, sodium methanolate, sodium ethanolate, potassium tert-butanolate, sodium acetate, potassium acetate, calcium acetate, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, potassium bicarbonate, sodium bicarbonate, or ammonium carbonate; and also tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dimethylbenzylamine pyridine, N-methylpiperidine, N-methylmorpholine, N,N-dimethylaminopyridine, diazabicyclooctane (DABCO), diazabicyclononene (DBN) or di-azabicycloundecene (DBU).

Suitable acid binders for carrying out the processes (b), (c), (d), (f), (g) and (h) according to the invention are in each case all inorganic and organic bases customary for such reactions. Preference is given to using alkaline earth metal or alkali metal hydrides, hydroxides, amides, alcoholates, acetates, fluorides, phosphates, carbonates or hydrogen carbonates, such as sodium hydride, sodium amide, lithium diisopropylamide, sodium methanolate, sodium ethanolate, potassium tert-butanolate, sodium

hydroxide, potassium hydroxide, sodium acetate, sodium phosphate, potassium phosphate, potassium fluoride, caesium fluoride, sodium carbonate, potassium carbonate, potassium hydrogencarbonate, sodium hydrogencarbonate or caesium carbonate; and also tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dimethyl-benzylamine, pyridine, N-methylpiperidine, N-methylmorpholine, N,N-dimethylaminopyridine, diazabicyclooctane (DABCO), diazabicyclononene (DBN) or diazabicycloundecene (DBU).

Suitable acid binders for carrying out the process (e) according to the invention are all inorganic and organic bases customary for such reactions. Preference is given to using alkaline earth metal or alkali metal hydrides, hydroxides, amides, alcoholates, acetates, carbonates or hydrogen carbonates, such as sodium hydride, sodium amide, lithium diisopropylamide, sodium methanolate, sodium ethanolate, potassium tert-butanolate, sodium acetate, potassium acetate, calcium acetate, ammonium acetate, sodium hydroxide, potassium hydroxide, ammonium hydroxide, sodium carbonate, potassium carbonate, potassium bicarbonate, sodium bicarbonate, or caesium carbonate; and also tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dimethylbenzylamine, pyridine, N-methylpiperidine, N-methylmorpholine, N,N-dimethylaminopyridine, diazabicyclooctane (DABCO), diazabicyclononene (DBN) or diazabicycloundecene (DBU).

Suitable condensing agents for carrying out the processes (a) and (i) according to the invention are all condensing agents customary for such amidation reactions. Preference is given to using acid halide former, such as phosgene, phosphorous tribromide, phosphorous trichloride, phosphorous pentachloride, phosphorous trichloride oxide or thionyl chloride; anhydride former, such as ethyl chloroformate, methyl chloroformate, isopropyl chloroformate, isobutyl chloroformate or methanesulfonyl chloride; carbodiimides, such as N,N'-dicyclohexylcarbodiimide (DCC) or other customary condensing agents, such as phosphorous pentoxide, polyphosphoric acid, N,N'-carbonyldiimidazole, 2-ethoxy-N-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), triphenylphosphine/tetrachloromethane or bromo-tripyrrolidinophosphonium-hexafluorophosphate.

The processes (a) and (i) according to the invention is optionally carried out in the presence of a catalyst. Preference is given to 4-dimethylaminopyridine, 1-hydroxy-benzotriazole or dimethylformamide.

The processes (b), (c), (d), (f), (g) and (h) according to the invention are carried out in the presence of a catalyst. Preference is given to palladium salts or complexes, such as palladium chloride, palladium acetate, tetrakis-(triphenylphosphine) palladium, bis-(triphenylphosphine) palladium dichloride or 1,1'-Bis(diphenylphosphino)ferrocenepalladium(II)chloride.



It is also possible to generate a palladium complex directly in the reaction mixture by separately adding to the reaction mixture a palladium salt and a complex ligand, such as triethylphosphane, tri-tert-butylphosphane, tricyclohexylphosphane, 2-(dicyclohexylphosphane)biphenyl, 2-(di-tert-butylphosphan)biphenyl, 2-(dicyclohexylphosphane)-2'-(N,N-dimethylamino)-biphenyl, triphenylphosphane, tris-(o-tolyl)phosphane, sodium 3-(diphenylphosphino)benzolsulfonate, tris-2-(methoxyphenyl)phosphane, 2,2'-bis-(diphenylphosphane)-1,1'-binaphthyl, 1,4-bis-(diphenylphosphane)butane, 1,2-bis-(diphenylphosphane)ethane, 1,4-bis-(dicyclohexylphosphane)butane, 1,2-bis-(dicyclohexylphosphane)ethane, 2-(dicyclohexylphosphane)-2'-(N,N-dimethylamino)-biphenyl, bis(diphenylphosphino)ferrocene or tris-(2,4-tert-butylphenyl)-phosphite.

10

When carrying out the processes (a) and (i) according to the invention, the reaction temperature can be varied within a relatively wide range. In general, the process is carried out at temperatures between 0°C and 150°C, preferably between 0°C and 120°C, particularly preferably between 10°C and 80°C.

15

When carrying out the processes (b), (c), (d), (f), (g) and (h) according to the invention, the reaction temperatures can in each case be varied within a relatively wide range. In general, the processes are carried out at temperatures between 0°C and 180°C, preferably between 10°C and 150°C, particularly preferably between 20°C and 120°C

20

When carrying out the process (e) according to the invention, the reaction temperature can be varied within a relatively wide range. In general, the process is carried out at temperatures between 0°C and 150°C, preferably between 20°C and 110°C.

25

When carrying out the process (a) according to the invention, in general between 0.8 and 15 mole, preferably of between 0.8 and 8 mole, of amine of the formula (III) and from 1 to 3 mole of acid binder are employed per mole of carboxylic acid derivative of the formula (II). However, it is also possible to employ the reaction components in other ratios. Work-up is carried out by customary methods. In general, water is added to the reaction mixture and the organic phase is separated off and, after drying, concentrated under reduced pressure. The residue that remains may, if appropriate, be freed of any impurities that may still be present using customary methods, such as chromatography or recrystallization.

30

When carrying out the process (b) according to the invention, in general 1 to 15 mole, preferably from 2 to 8 mole, of boronic acid derivative of the formula (V) and from 1 to 5 mol of acid binder are employed per mole of halogeno-carboxamide of the formula (IV). However, it is also possible to employ the reaction components in other ratios. Work-up is carried out by customary methods. In general, water is added to the reaction mixture and the precipitate is separated off and dried. The

35

residue that remains may, if appropriate, be freed of any impurities that may still be present using customary methods, such as chromatography or recrystallization.

When carrying out the process (c) according to the invention, in general 0.8 to 15 mole, preferably from 0.8 to 8 mole, of pyridinyl derivative of the formula (VII) and from 1 to 10 mol of acid binder and from 0.5 to 5 mole% of a catalyst are employed per mole of carboxamide boronic acid derivative of the formula (VI). However, it is also possible to employ the reaction components in other ratios. Work-up is carried out by customary methods. In general, water is added to the reaction mixture and the precipitate is separated off and dried. The residue that remains may, if appropriate, be freed of any impurities that may still be present using customary methods, such as chromatography or recrystallization.

When carrying out the process (d) according to the invention, in general 0.8 to 15 mole, preferably from 0.8 to 8 mole, of pyridinyl derivative of the formula (VII) and from 0.8 to 15 mole, preferably from 0.8 to 8 mole, of 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bis-1,3,2-dioxaborolane and from 1 to 5 mol of acid binder and from 1 to 5 mol of a catalyst are employed per mole of carboxamide derivative of the formula (IV). However, it is also possible to employ the reaction components in other ratios. Work-up is carried out by customary methods. In general, water is added to the reaction mixture and the precipitate is separated off and dried. The residue that remains may, if appropriate, be freed of any impurities that may still be present using customary methods, such as chromatography or recrystallization.

When carrying out process (e) according to the invention, per mole of the pyridinylanilide of the formula (I-1) in general 0.2 to 5 mole, preferably 0.5 to 2 mole of an halogenide of the formula (VIII) are employed. However, it is also possible to employ the reaction components in other ratios. Work-up is carried out by customary methods.

All processes according to the invention are generally each carried out under atmospheric pressure. However, in each case it is also possible to operate under elevated or reduced pressure – in general between 0,1 bar and 10 bar.

30

The substances according to the invention have potent microbicidal activity and can be employed for controlling unwanted micro-organisms, such as fungi and bacteria, in crop protection and in the protection of materials.

Fungicides can be employed in crop protection for controlling Plasmodiophoromycetes, Oomycetes, Chytridiomycetes, Zygomycetes, Ascomycetes, Basidiomycetes and Deuteromycetes.

Bactericides can be employed in crop protection for controlling Pseudomonadaceae, Rhizobiaceae, Enterobacteriaceae, Corynebacteriaceae and Streptomycetaceae.

- Some pathogens causing fungal and bacterial diseases which come under the generic names listed above may be mentioned as examples, but not by way of limitation:
- 5 Xanthomonas species, such as, for example, Xanthomonas campestris pv. oryzae;  
Pseudomonas species, such as, for example, Pseudomonas syringae pv. lachrymans;  
Erwinia species, such as, for example, Erwinia amylovora;  
Pythium species, such as, for example, Pythium ultimum;
- 10 Phytophthora species, such as, for example, Phytophthora infestans;  
Pseudoperonospora species, such as, for example, Pseudoperonospora humuli or Pseudoperonospora cubensis;  
Plasmopara species, such as, for example, Plasmopara viticola;  
Bremia species, such as, for example, Bremia lactucae;
- 15 Peronospora species, such as, for example, Peronospora pisi or P. brassicae;  
Erysiphe species, such as, for example, Erysiphe graminis;  
Sphaerotheca species, such as, for example, Sphaerotheca fuliginea;  
Podosphaera species, such as, for example, Podosphaera leucotricha;  
Venturia species, such as, for example, Venturia inaequalis;
- 20 Pyrenophora species, such as, for example, Pyrenophora teres or P. graminea  
(conidia form: Drechslera, syn: Helminthosporium);  
Cochliobolus species, such as, for example, Cochliobolus sativus  
(conidia form: Drechslera, syn: Helminthosporium);  
Uromyces species, such as, for example, Uromyces appendiculatus;
- 25 Puccinia species, such as, for example, Puccinia recondita;  
Sclerotinia species, such as, for example, Sclerotinia sclerotiorum;  
Tilletia species, such as, for example, Tilletia caries;  
Ustilago species, such as, for example, Ustilago nuda or Ustilago avenae;  
Pellicularia species, such as, for example, Pellicularia sasakii;
- 30 Pyricularia species, such as, for example, Pyricularia oryzae;  
Fusarium species, such as, for example, Fusarium culmorum;  
Botrytis species, such as, for example, Botrytis cinerea;  
Septoria species, such as, for example, Septoria nodorum;  
Leptosphaeria species, such as, for example, Leptosphaeria nodorum;
- 35 Cercospora species, such as, for example, Cercospora canescens;  
Alternaria species, such as, for example, Alternaria brassicae; and

Pseudocercospora species, such as, for example, Pseudocercospora herpotrichoides.

The active compounds according to the invention also have very good fortifying action in plants. Accordingly, they can be used for mobilizing the defences of the plant against attack by unwanted  
5 micro-organisms.

In the present context, plant-fortifying (resistance-inducing) substances are to be understood as meaning those substances which are capable of stimulating the defence system of plants such that, when the treated plants are subsequently inoculated with unwanted micro-organisms, they show  
10 substantial resistance against these micro-organisms.

In the present case, unwanted micro-organisms are to be understood as meaning phytopathogenic fungi, bacteria and viruses. Accordingly, the substances according to the invention can be used to protect plants for a certain period after the treatment against attack by the pathogens mentioned. The  
15 period for which protection is provided generally extends over 1 to 10 days, preferably 1 to 7 days, after the treatment of the plants with the active compounds.

The fact that the active compounds are well tolerated by plants at the concentrations required for controlling plant diseases permits the treatment of above-ground parts of plants, of propagation stock  
20 and seeds, and of the soil.

The active compounds according to the invention are also suitable for increasing the yield of crops. In addition, they show reduced toxicity and are well tolerated by plants.

25 At certain concentrations and application rates, the active compounds according to the invention can also be used as herbicides, for influencing plant growth and for controlling animal pests. They can also be used as intermediates and precursors for the synthesis of further active compounds.

The active compounds according to the invention can be used to treat all plants and parts of plants.  
30 By plants are understood here all plants and plant populations such as desired and undesired wild plants or crop plants (including naturally occurring crop plants). Crop plants can be plants which can be obtained by conventional breeding and optimization methods or by biotechnological and genetic engineering methods or combinations of these methods, including the transgenic plants and including the plant varieties which can or cannot be protected by plant varieties property rights. Parts of plants  
35 are to be understood as meaning all above-ground and below-ground parts and organs of plants, such as shoot, leaf, flower and root, examples which may be mentioned being leaves, needles, stems,

trunks, flowers, fruit-bodies, fruits and seeds and also roots, tubers and rhizomes. Parts of plants also include harvested plants and vegetative and generative propagation material, for example seedlings, tubers, rhizomes, cuttings and seeds.

- 5 The treatment of the plants and the parts of plants with the active compounds according to the invention is carried out directly or by action on their surroundings, habitat or storage space, according to customary treatment methods, for example by dipping, spraying, evaporating, atomizing, broadcasting, spreading-on and, in the case of propagation material, in particular in the case of seeds, furthermore by one- or multi-layer coating.

10

In the protection of materials, the compounds according to the invention can be employed for protecting industrial materials against infection with, and destruction by, undesired micro-organisms.

- 15 Industrial materials in the present context are understood as meaning non-living materials which have been prepared for use in industry. For example, industrial materials which are intended to be protected by active compounds according to the invention from microbial change or destruction can be adhesives, sizes, paper and board, textiles, leather, wood, paints and plastic articles, cooling lubricants and other materials which can be infected with, or destroyed by, micro-organisms. Parts of production plants, for example cooling-water circuits, which may be impaired by the proliferation of
- 20 micro-organisms may also be mentioned within the scope of the materials to be protected. Industrial materials which may be mentioned within the scope of the present invention are preferably adhesives, sizes, paper and board, leather, wood, paints, cooling lubricants and heat-transfer liquids, particularly preferably wood.

- 25 Micro-organisms capable of degrading or changing the industrial materials which may be mentioned are, for example, bacteria, fungi, yeasts, algae and slime organisms. The active compounds according to the invention preferably act against fungi, in particular moulds, wood-discolouring and wood-destroying fungi (Basidiomycetes), and against slime organisms and algae.

- 30 Micro-organisms of the following genera may be mentioned as examples:

*Alternaria*, such as *Alternaria tenuis*,

*Aspergillus*, such as *Aspergillus niger*,

*Chaetomium*, such as *Chaetomium globosum*,

*Coniophora*, such as *Coniophora puetana*,

- 35 *Lentinus*, such as *Lentinus tigrinus*,

*Penicillium*, such as *Penicillium glaucum*,

- Polyporus, such as *Polyporus versicolor*,  
Aureobasidium, such as *Aureobasidium pullulans*,  
Sclerophoma, such as *Sclerophoma pityophila*,  
Trichoderma, such as *Trichoderma viride*,  
5 *Escherichia*, such as *Escherichia coli*,  
*Pseudomonas*, such as *Pseudomonas aeruginosa*, and  
*Staphylococcus*, such as *Staphylococcus aureus*.

Depending on their particular physical and/or chemical properties, the active compounds can be  
10 converted to the customary formulations, such as solutions, emulsions, suspensions, powders, foams,  
pastes, granules, aerosols and microencapsulations in polymeric substances and in coating  
compositions for seeds, and ULV cool and warm fogging formulations.

These formulations are produced in a known manner, for example by mixing the active compounds  
15 with extenders, that is, liquid solvents, liquefied gases under pressure, and/or solid carriers, optionally  
with the use of surfactants, that is emulsifiers and/or dispersants, and/or foam formers. If the extender  
used is water, it is also possible to employ, for example, organic solvents as auxiliary solvents.  
Essentially, suitable liquid solvents are: aromatics such as xylene, toluene or alkylnaphthalenes,  
chlorinated aromatics or chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes  
20 or methylene chloride, aliphatic hydrocarbons such as cyclohexane or paraffins, for example  
petroleum fractions, alcohols such as butanol or glycol and their ethers and esters, ketones such as  
acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents such  
as dimethylformamide or dimethyl sulphoxide, or else water. Liquefied gaseous extenders or carriers  
are to be understood as meaning liquids which are gaseous at standard temperature and under  
25 atmospheric pressure, for example aerosol propellants such as halogenated hydrocarbons, or else  
butane, propane, nitrogen and carbon dioxide. Suitable solid carriers are: for example ground natural  
minerals such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous  
earth, and ground synthetic minerals such as finely divided silica, alumina and silicates. Suitable solid  
carriers for granules are: for example crushed and fractionated natural rocks such as calcite, marble,  
30 pumice, sepiolite and dolomite, or else synthetic granules of inorganic and organic meals, and  
granules of organic material such as sawdust, coconut shells, maize cobs and tobacco stalks. Suitable  
emulsifiers and/or foam formers are: for example nonionic and anionic emulsifiers, such as  
polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl  
polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates, or else protein hydrolysates.  
35 Suitable dispersants are: for example lagnosulphite waste liquors and methylcellulose.

Tackifiers such as carboxymethylcellulose and natural and synthetic polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, or else natural phospholipids such as cephalins and lecithins and synthetic phospholipids can be used in the formulations. Other possible additives are mineral and vegetable oils.

5

It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic dyestuffs such as alizarin dyestuffs, azo dyestuffs and metal phthalocyanine dyestuffs, and trace nutrients such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

10

The formulations generally comprise between 0.1 and 95 percent by weight of active compound, preferably between 0.5 and 90%.

The active compounds according to the invention can be used as such or in their formulations, also in a mixture with known fungicides, bactericides, acaricides, nematocides or insecticides, to broaden, for example, the activity spectrum or to prevent development of resistance. In many cases, synergistic effects are obtained, i.e. the activity of the mixture is greater than the activity of the individual components.

20 Examples of suitable mixing components are the following:

**Fungicides:**

2-phenylphenol; 8-hydroxychinolinsulfat; acibenzolar-S-methyl; aldimorph; amidoflumet; ampropylfos; ampropylfos-potassium; andoprim; anilazine; azaconazole; azoxystrobin; benalaxyl; benodanil; benomyl; benthiavalicarb-isopropyl; benzamacril; benzamacril-isobutyl; bilanafos; binapacryl; biphenyl; 25 bitertanol; blasticidin-S; bromuconazole; bupirimate; buthiobate; butylamin; calcium polysulfide; capsimycin; captafol; captan; carbendazim; carboxin; carpropamid; carvone; chinomethionat; chlorthiazole; chlorfenazole; chloroneb; chlorothalonil; chlozolinate; clozylacon; cyazofamid; cyflufenamid; cymoxanil; cyproconazole; cyprodinil; cyprofuram; Dagger G; debacarb; dichlofluanid; dichlone; dichlorophen; diclocymet; diclomezine; dicloran; diethofencarb; difenoconazole; diflumetorim; dimethirimol; dimethomorph; dimoxystrobin; diniconazole; diniconazole-M; dinocap; diphenylamine; dipyrithione; ditalimfos; dithianon; dodine; drazoxolon; edifenphos; epoxiconazole; ethaboxam; ethirimol; etridiazole; famoxadone; fenamidone; fenapanil; fenarimol; fenbuconazole; fenfuram; fenhexamid; fenitropan; fenoxanil; fenpiclonil; fenpropidin; fenpropimorph; ferbam; fluazinam; flubenzimine; fludioxonil; flumetover; flumorph; fluoromide; fluoxastrobin; fluquinconazole; flurprimidol; flusilazole; 35 flusulfamide; flutolanil; flutriafol; folpet; fosetyl-Al; fosetyl-sodium; fuberidazole; furalaxyl; furametpyr; furcarbanil; furmecyclox; guazatine; hexachlorobenzene; hexaconazole; hymexazol; imazalil;

imibenconazole; iminoctadine triacetate; iminoctadine tris(albesil; iodocarb; ipconazole; iprobenfos;  
 iprodione; iprovalicarb; irumamycin; isoprothiolane; isovaledione; kasugamycin; kresoxim-methyl;  
 mancozeb; maneb; meferimzone; mepanipyrim; mepronil; metalaxyl; metalaxyl-m; metconazole; metha-  
 sulfocarb; methfuroxam; metiram; metominostrobin; metsulfovax; mildiomyacin; myclobutanil; myclo-  
 5 zolin; natamycin; nicobifen; nitrothal-isopropyl; noviflumuron; nuarimol; ofurace; orysastrobin;  
 oxadixyl; oxolinic acid; oxpoconazole; oxycarboxin; oxyfenthiiin; paclobutrazol; pefurazoate; pencon-  
 azole; pencycuron; phosdiphen; phthalide; picoxystrobin; piperalin; polyoxins; polyoxorim; proben-  
 azole; prochloraz; procymidone; propamocarb; propanosine-sodium; propiconazole; propineb; pro-  
 quinazid; prothioconazole; pyraclostrobin; pyrazophos; pyrifenoxy; pyrimethanil; pyroquilon; pyroxyfur;  
 10 pyrrolnitrine; quinconazole; quinoxifen; quintozone; simeconazole; spiroxamine; sulfur; tebuconazole;  
 tecloftalam; tecnazene; tetcyclacis; tetraconazole; thiabendazole; thicyofen; thifluzamide; thiophanate-  
 methyl; thiram; tioxyimid; tolclofos-methyl; tolylfluanid; triadimefon; triadimenol; triazbutil; triazoxide;  
 tricyclamide; tricyclazole; tridemorph; trifloxystrobin; triflumizole; triforine; triticonazole; uniconazole;  
 validamycin a; vinclozolin; zineb; ziram; zoxamide; (2S)-N-[2-[4-[[3-(4-chlorophenyl)-2-propynyl]oxy]-  
 15 3-methoxyphenyl]ethyl]-3-methyl-2-[(methylsulphonyl)amino]butanamide; 1-(1-naphthalenyl)-1H-pyr-  
 role-2,5-dione; 2,3,5,6-tetrachloro-4-(methylsulphonyl)pyridine; 2-amino-4-methyl-N-phenyl-5-thiazole-  
 carboxamide; 2-chloro-N-(2,3-dihydro-1,1,3-trimethyl-1H-inden-4-yl)-3-pyridinecarboxamide; 3,4,5-  
 trichloro-2,6-pyridinedicarbonitrile; actinovate; cis-1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cyclo-  
 heptanol; methyl 1-(2,3-dihydro-2,2-dimethyl-1H-inden-1-yl)-1H-imidazole-5-carboxylate; monopotas-  
 20 sium carbonate; N-(6-methoxy-3-pyridinyl)-cyclopropanecarboxamide; N-butyl-8-(1,1-dimethylethyl)-1-  
 oxaspiro[4.5]decane-3-amine; sodium tetrathiocarbonate; and copper salts and preparations, such as  
 Bordeaux mixture; copper hydroxide; copper naphthenate; copper oxychloride; copper sulphate;  
 cufraneb; copper oxide; mancopper; oxine-copper.

## 25 Bactericides:

bronopol, dichlorophen, nitrapyrin, nickel dimethyldithiocarbamate, kasugamycin, octhilineone,  
 furancarboxylic acid, oxytetracyclin, probenazole, streptomycin, tecloftalam, copper sulphate and  
 other copper preparations.

## 30 Insecticides / acaricides / nematocides:

abamectin, ABG-9008, acephate, acequinocyl, acetamiprid, acetoprole, acrinathrin, AKD-1022, AKD-  
 3059, AKD-3088, alanycarb, aldicarb, aldoxycarb, allethrin, allethrin 1R-isomers, alpha-cypermethrin  
 (alphamethrin), amidoflumet, aminocarb, amitraz, avermectin, AZ-60541, azadirachtin, azamethphos,  
 azinphos-methyl, azinphos-ethyl, azocyclotin, Bacillus popilliae, Bacillus sphaericus, Bacillus subtilis,  
 35 Bacillus thuringiensis, Bacillus thuringiensis strain EG-2348, Bacillus thuringiensis strain GC-91,  
 Bacillus thuringiensis strain NCTC-11821, baculoviruses, Beauveria bassiana, Beauveria tenella, ben-



diocarb, benfuracarb, bensultap, benzoximate, beta-cyfluthrin, beta-cypermethrin, bifenazate, bifenthrin,  
 binapacryl, bioallethrin, bioallethrin-S-cyclopentyl-isomer, bioethanomethrin, biopermethrin, biores-  
 methrin, bistrifluron, BPMC, brofenprox, bromophos-ethyl, bromopropylate, bromfeninfos (-methyl),  
 BTG-504, BTG-505, bufencarb, buprofezin, butathiofos, butocarboxim, butoxycarboxim, butyl-  
 5 pyridaben, cadusafos, camphechlor, carbaryl, carbofuran, carbophenothion, carbosulphan, cartap, CGA-  
 50439, chinomethionat, chlordane, chlordimeform, chloethocarb, chlorethoxyfos, chlorfenapyr, chlor-  
 fenvinphos, chlorfluazuron, chlormephos, chlorobenzilate, chloropicrin, chlorproxyfen, chlorpyrifos-  
 methyl, chlorpyrifos (-ethyl), chlovaporthrin, chromafenozide, cis-cypermethrin, cis-resmethrin, cis-per-  
 methrin, clocythrln, cloethocarb, clofentezine, clothianidin, clothiazoben, codlemone, coumaphos,  
 10 cyanofenphos, cyanophos, cycloprene, cycloprothrin, Cydia pomonella, cyfluthrin, cyhalothrin, cyhexa-  
 tin, cypermethrin, cyphenothrin (1R-trans-isomer), cyromazine, DDT, deltamethrin, demeton-S-methyl,  
 demeton-S-methylsulphone, diafenthiuron, dialifos, diazinon, dichlofenthion, dichlorvos, dicofol, di-  
 crotophos, dicyclanil, diflubenzuron, dimethoate, dimethylvinphos, dinobuton, dinocap, dinotefuran,  
 diofenolan, disulphoton, docusat-sodium, dofenapyn, DOWCO-439, eflusilanate, emamectin, ema-  
 15 mectin-benzoate, empenethrin (1R-isomer), endosulphan, Entomophthora spp., EPN, esfenvalerate, ethio-  
 fencarb, ethiprole, ethion, ethoprophos, etofenprox, etoxazole, etrimfos, famphur, fenamiphos, fenaza-  
 quin, fenbutatin oxide, fenfluthrin, fenitrothion, fenobucarb, fenothiocab, fenoxacrim, fenoxycarb, fen-  
 propathrin, fenpyrad, fenpyrithrin, fenpyroximate, fensulphothion, fenthion, fentrifanil, fenvalerate,  
 fipronil, flonicamid, fluacrypyrim, fluazuron, flubenzimine, flubrocylthrin, flucycloxuron, flu-  
 20 cythrin, flufenerim, flufenoxuron, flufenprox, flumethrin, flupyrzofos, flutenzin (flufenzine), flu-  
 valinate, fonofos, formetanate, formothion, fosmethilan, fosthiazate, fubfenprox (fluproxyfen), furathio-  
 carb, gamma-HCH, gossypure, grandlure, granulosis viruses, halfenprox, halofenozide, HCH, HCN-  
 801, heptenophos, hexaflumuron, hexythiazox, hydramethylnone, hydroprene, IKA-2002, imidacloprid,  
 imiprothrin, indoxacarb, iodofenphos, iprobenfos, isazofos, isofenphos, isoprocarb, isoxathion,  
 25 ivermectin, japonilure, kadethrin, nuclear polyhedrosis viruses, kinoprene, lambda-cyhalothrin, lindane,  
 lufenuron, malathion, mecarbarn, mesulphenfos, metaldehyde, metam-sodium, methacrifos, methamido-  
 phos, Metharhizium anisopliae, Metharhizium flavoviride, methidathion, methiocarb, methomyl, metho-  
 prene, methoxychlor, methoxyfenozide, metolcarb, metoxadiazone, mevinphos, milbemectin, milbe-  
 mycin, MKI-245, MON-45700, monocrotophos, moxidectin, MTI-800, naled, NC-104, NC-170, NC-  
 30 184, NC-194, NC-196, niclosamide, nicotine, nitenpyram, nithiazine, NNI-0001, NNI-0101, NNI-0250,  
 NNI-9768, novaluron, noviflumuron, OK-5101, OK-5201, OK-9601, OK-9602, OK-9701, OK-9802,  
 omethoate, oxamyl, oxydemeton-methyl, Paecilomyces fumosoroseus, parathion-methyl, parathion (-  
 ethyl), permethrin (cis-, trans-), petroleum, PH-6045, phenothrin (1R-trans isomer), phenthoate, phorate,  
 phosalone, phosmet, phosphamidon, phosphocarb, phoxim, piperonyl butoxide, pirimicarb, pirimiphos-  
 35 methyl, pirimiphos-ethyl, prallethrin, profenofos, promecarb, propaphos, propargite, propetamphos, pro-  
 poxur, prothiofos, prothoate, protrifenbutate, pymetrozine, pyraclofos, pyresmethrin, pyrethrum,

pyridaben, pyridalyl, pyridaphenthion, pyridathion, pyrimidifen, pyriproxyfen, quinalphos, resmethrin, RH-5849, ribavirin, RU-12457, RU-15525, S-421, S-1833, salithion, sebufos, SI-0009, silafluofen, spinosad, spirodiclofen, spiromesifen, sulphluramid, sulphotep, sulprofos, SZI-121, tau-fluvalinate, tebufenozide, tebufenpyrad, tebupirimfos, teflubenzuron, tefluthrin, temephos, temvinphos, terbam, terbufos, tetrachlorvinphos, tetradifon, tetramethrin, tetramethrin (1R-isomer), tetrasul, theta-cypermethrin, thiocloprid, thiamethoxam, thiapronil, thiatriphos, thiocyclam hydrogenoxalate, thiodicarb, thiofanox, thiometon, thiosultap-sodium, thuringiensin, tolfenpyrad, tralocythrin, tralomethrin, transfluthrin, triarathene, triazamate, triazophos, triazuron, trichlophenidine, trichlorfon, triflumuron, trimethacarb, vamidothion, vaniliprole, verbutin, *Verticillium lecanii*, WL-108477, WL-40027, YI-5201, YI-5301, YI-5302, XMC, xylilcarb, ZA-3274, zeta-cypermethrin, zolaprofos, ZXI-8901, the compound 3-methylphenyl propylcarbamate (tsunacide Z), the compound 3-(5-chloro-3-pyridinyl)-8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octane-3-carbonitrile (CAS-Reg. No. 185982-80-3) and the corresponding 3-endo-isomer (CAS-Reg. No. 185984-60-5) (cf. WO-96/37494, WO-98/25923), and preparations which comprise insecticidally active plant extracts, nematodes, fungi or viruses.

15

A mixture with other known active compounds, such as herbicides, or with fertilizers and growth regulators, safeners and/or semiochemicals is also possible.

20

In addition, the compounds of the formula (I) according to the invention also have very good antimycotic activity. They have a very broad antimycotic activity spectrum in particular against dermatophytes and yeasts, moulds and diphasic fungi (for example against *Candida* species such as *Candida albicans*, *Candida glabrata*) and *Epidermophyton floccosum*, *Aspergillus* species such as *Aspergillus niger* and *Aspergillus fumigatus*, *Trichophyton* species such as *Trichophyton mentagrophytes*, *Microsporon* species such as *Microsporon canis* and *audouinii*. The list of these

25

fungi does by no means limit the mycotic spectrum which can be covered, but is only for illustration.

The active compounds can be used as such, in the form of their formulations or the use forms prepared therefrom, such as ready-to-use solutions, suspensions, wettable powders, pastes, soluble powders, dusts and granules. Application is carried out in a customary manner, for example by watering, spraying, atomizing, broadcasting, dusting, foaming, spreading, etc. It is furthermore possible to apply the active compounds by the ultra-low volume method, or to inject the active compound preparation or the active compound itself into the soil. It is also possible to treat the seeds of the plants.

30

When using the active compounds according to the invention as fungicides, the application rates can be varied within a relatively wide range, depending on the kind of application. For the treatment of parts of plants, the active compound application rates are generally between 0.1 and 10,000 g/ha,

35

preferably between 10 and 1000 g/ha. For seed dressing, the active compound application rates are generally between 0.001 and 50 g per kilogram of seed, preferably between 0.01 and 10 g per kilogram of seed. For the treatment of the soil, the active compound application rates are generally between 0.1 and 10,000 g/ha, preferably between 1 and 5,000 g/ha.

5

As already mentioned above, it is possible to treat all plants and their parts according to the invention. In a preferred embodiment, wild plant species and plant cultivars, or those obtained by conventional biological breeding, such as crossing or protoplast fusion, and parts thereof, are treated. In a further preferred embodiment, transgenic plants and plant cultivars obtained by genetic engineering, if appropriate in combination with conventional methods (Genetically Modified Organisms), and parts thereof, are treated. The term "parts" or "parts of plants" or "plant parts" has been explained above.

10

Particularly preferably, plants of the plant cultivars which are in each case commercially available or in use are treated according to the invention. Plant cultivars are to be understood as meaning plants having new properties ("traits") and which have been obtained by conventional breeding, by mutagenesis or by recombinant DNA techniques. They can be cultivars, varieties, bio- or genotypes.

15

Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive ("synergistic") effects. Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity of the substances and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products are possible which exceed the effects which were actually to be expected.

20

25

The transgenic plants or plant cultivars (i.e. those obtained by genetic engineering) which are preferably to be treated according to the invention include all plants which, in the genetic modification, received genetic material which imparted particularly advantageous useful properties ("traits") to these plants. Examples of such properties are better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products. Further and particularly emphasized examples of such properties are a better defence of the plants against animal and microbial pests, such as against insects, mites, phytopathogenic fungi, bacteria and/or


30

35

viruses, and also increased tolerance of the plants to certain herbicidally active compounds. Examples of transgenic plants which may be mentioned are the important crop plants, such as cereals (wheat, rice), maize, soya beans, potatoes, cotton, tobacco, oilseed rape and also fruit plants (with the fruits apples, pears, citrus fruits and grapes), and particular emphasis is given to maize, soya beans, potatoes, cotton, tobacco and oilseed rape. Traits that are emphasized are in particular increased defence of the plants against insects by toxins formed in the plants, in particular those formed in the plants by the genetic material from *Bacillus thuringiensis* (for example by the genes CryIA(a), CryIA(b), CryIA(c), CryIIA, CryIIIA, CryIIIB2, Cry9c, Cry2Ab, Cry3Bb and CryIF and also combinations thereof) (hereinbelow referred to as "Bt plants"). Traits that are also particularly emphasized are the increased defence of the plants against fungi, bacteria and viruses by systemic acquired resistance (SAR), systemin, phytoalexins, elicitors and resistance genes and correspondingly expressed proteins and toxins. Traits that are furthermore particularly emphasized are the increased tolerance of the plants to certain herbicidally active compounds, for example imidazolinones, sulphonylureas, glyphosate or phosphinotricin (for example the "PAT" gene). The genes which impart the desired traits in question can also be present in combination with one another in the transgenic plants. Examples of "Bt plants" which may be mentioned are maize varieties, cotton varieties, soya bean varieties and potato varieties which are sold under the trade names YIELD GARD® (for example maize, cotton, soya beans), KnockOut® (for example maize), StarLink® (for example maize), Bollgard® (cotton), Nucoton® (cotton) and NewLeaf® (potato). Examples of herbicide-tolerant plants which may be mentioned are maize varieties, cotton varieties and soya bean varieties which are sold under the trade names Roundup Ready® (tolerance to glyphosate, for example maize, cotton, soya bean), Liberty Link® (tolerance to phosphinotricin, for example oilseed rape), IMI® (tolerance to imidazolinones) and STS® (tolerance to sulphonylureas, for example maize). Herbicide-resistant plants (plants bred in a conventional manner for herbicide tolerance) which may be mentioned also include the varieties sold under the name Clearfield® (for example maize). Of course, these statements also apply to plant cultivars which have these genetic traits or genetic traits still to be developed, and which will be developed and/or marketed in the future.

The plants listed can be treated according to the invention in a particularly advantageous manner with the compounds of the general formula (I) or the active compound mixtures according to the invention. The preferred ranges stated above for the active compounds or mixtures also apply to the treatment of these plants. Particular emphasis is given to the treatment of plants with the compounds or mixtures specifically mentioned in the present text.


The preparation and the use of the active compounds according to the invention is illustrated by the examples below.

**Preparation examples****Example 1**

(I-8-1)

- 5 A solution of 10.0 g (25.6 mmol) *N*-(2-iodophenyl)-2-(trifluoromethyl)benzamide, 8.0 g (31.5 mmol) bis(pinacolato)diboron, 7.4 g (75.4 mmol) potassium acetate and 0.18 g (0.25 mmol) 1,1'-bis-(diphenylphosphino)ferrocenepalladium(II)chloride in 120 ml dimethyl sulphoxide was heated under an inert gas atmosphere for 2 h at 90°C. At room temperature 6.5 g (25.0 mmol) 2-bromo-3-chloro-5-(trifluoromethyl)pyridine, 70.0 ml of a 2 M sodium carbonate solution and 0.18 g (0.25 mmol) 1,1'-
- 10 bis(diphenylphosphino)ferrocenepalladium(II)chloride were added. The reaction mixture was heated for 16 h at 90°C. For work-up the mixture was poured into water, extracted with ethyl acetate, dried over sodium sulphate, filtrated and concentrated in vacuo. Column chromatography (cyclohexane/ethyl acetate 3:1) yielded 1.5 g (3.4 mmol, 13 %) of *N*-(2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]phenyl)-2-(trifluoromethyl)benzamide [log P (pH 2.3) = 3.58; compound No. I-8-1 in Table 3].

15

**Example 2**


(I-8-15)

- A mixture of 0.2 g (0.8 mmol) of 6-(2-amino-phenyl)-pyridine-3-carbaldehyde (*E*)-*O*-methyl-oxime (III-2), 0.155 mg (0.88 mmol) of 2-chloronicotiny chloride, 122 mg (0.88 mmol) of potassium carbonate in 20 ml of acetonitrile was stirred for 15 hours at room temperature. For the work-up, the reaction mixture was poured into water, extracted with ethyl acetate, dried over sodium sulphate and concentrated under vacuum. The solid was triturated with petroleum ether and filtered, to give
- 20



150 mg (0.41 mmol, 41 %) of 2-chloro-*N*-(2-[5-((*E*)methoxyimino-methyl)-pyridin-2-yl]-phenyl)-nicotinamide as a white powder [ $\log P$  (pH 2.3) = 2.95; compound No. I-8-15 in Table 3].

The pyridinylanilides of the formula (I) listed in the following tables below are likewise prepared analogously to Examples 1 and 2 described above and in accordance with the general descriptions of the processes.


**Table 1**







No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R	A	logP
I-6-1	2-F	H	H	H		2.37
I-6-2	H	H	H	3-CH <sub>3</sub>		2.83
I-6-3	H	H	H	H		2.43
I-6-4	6-Cl	H	H	H		2.00
I-6-5	6-Cl	H	H	H		2.82
I-6-6	6-Cl	H	H	H		2.74
I-6-7	6-Cl	H	H	H		2.10


No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R	A	logP
I-6-8	6-Cl	H	H	H		2.00
I-6-9	6-Cl	H	H	H		2.64

**Table 2**







No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R	A	logP
I-7-1	H	H	H	4-CF <sub>3</sub>		3.63
I-7-2	H	H	H	H		
I-7-3	2-Br	H	H	H		2.04
I-7-4	2-Br	H	H	H		2.02
I-7-5	2-Br	H	H	H		2.85
I-7-6	2-Br	H	H	H		2.13












No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R	A	logP
I-7-7	2-Cl	H	H	H		1.97
I-7-8	2-Cl	H	H	H		2.60
I-7-9	2-Cl	6-Cl	H	H		2.57
I-7-10	2-Cl	6-Cl	H	H		3.27













**Table 3**












(I-8)













No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R	A	logP m.p./°C
I-8-1	3-Cl	5-CF <sub>3</sub>	H	H		3.59 139
I-8-2	3-Cl	5-CF <sub>3</sub>	H	H		3.44
I-8-3	3-Cl	5-CF <sub>3</sub>	H	H		3.83
I-8-4	4-CH <sub>3</sub>	5-CH=CH-CH=CH-6		H		2.79








No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R	A	logP m.p./°C
I-8-5	4-CH <sub>3</sub>	5-CH=CH-CH=CH-6		H		2.39
I-8-6	4-CH <sub>3</sub>	5-CH=CH-CH=CH-6		H		3.97
I-8-7	H	H	H	5-Cl		3.49
I-8-8	H	H	H	3-CH <sub>3</sub>		2.86
I-8-9	5-CF <sub>3</sub>	H	H	H		4.13
I-8-10	5-Cl	H	H	H		2.95
I-8-11	5-Cl	H	H	H		3.06
I-8-12	5-Cl	H	H	H		4.00
I-8-13	5-Cl	H	H	H		4.13
I-8-14	5-Cl	H	H	H		3.05
I-8-15	5-CH=N-OCH <sub>3</sub>	H	H	H		2.95

No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R	A	logP m.p./°C
I-8-16	5-CH=N-OCH <sub>3</sub>	H	H	H		3.03
I-8-17	5-CH=N-OCH <sub>3</sub>	H	H	H		3.93
I-8-18	5-CH=N-OCH <sub>3</sub>	H	H	H		4.00
I-8-19	5-CH=N-OCH <sub>3</sub>	H	H	H		3.07
I-8-20	3-Cl	5-CF <sub>3</sub>	H	H		3.45
I-8-21	3-Cl	5-CF <sub>3</sub>	H	H		3.49
I-8-22	3-Cl	5-CF <sub>3</sub>	H	H		3.61
I-8-23	3-Cl	5-CF <sub>3</sub>	H	H		3.04
I-8-24	3-Cl	5-CF <sub>3</sub>	H	H		3.59
I-8-25	3-Cl	5-CF <sub>3</sub>	H	H		3.43
I-8-26	3-Cl	5-CF <sub>3</sub>	H	H		3.55
I-8-27	3-Cl	5-CF <sub>3</sub>	H	H		2.91


No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R	A	logP m.p./°C
I-8-28	3-Cl	5-CF <sub>3</sub>	H	H		3.48
I-8-29	3-Cl	5-CF <sub>3</sub>	H	H		3.13
I-8-30	3-Cl	5-CF <sub>3</sub>	H	H		2.92
I-8-31	3-Cl	5-CF <sub>3</sub>	H	H		3.29
I-8-32	5-CH <sub>3</sub>	H	H	H		1.68
I-8-33	5-CH <sub>3</sub>	H	H	H		2.67
I-8-34	3-Cl	5-Cl	H	H		2.81
I-8-35	3-Cl	5-Cl	H	H		3.75
I-8-36	3-Cl	5-Cl	H	H		3.40
I-8-37	3-Cl	5-Cl	H	H		2.53
I-8-38	3-Cl	5-Cl	H	H		2.93

No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R	A	logP m.p./°C
I-8-39	5-Br	H	H	H		3.95
I-8-40	5-Br	H	H	H		4.47
I-8-41	3-Cl	5-Cl	H	H		3.30
I-8-42	3-Cl	5-Cl	H	H		3.45
I-8-43	3-Cl	5-Cl	H	H		3.83
I-8-44	3-Cl	5-Cl	H	H		3.75
I-8-45	3-Cl	5-Cl	H	H		3.19
I-8-46	3-Cl	5-Cl	H	H		3.27
I-8-47	5-CH <sub>3</sub>	H	H	H		3.20
I-8-48	3-CH <sub>3</sub>	5-Br	H	H		3.21
I-8-49	3-Cl	5-Br	H	H		3.40
I-8-50	3-Cl	5-Br	H	H		3.27

No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R	A	logP m.p./°C
I-8-51	3-CH <sub>3</sub>	5-Br	H	H		3.30
I-8-52	3-CH <sub>3</sub>	5-Br	H	H		2.71
I-8-53	3-Cl	5-Br	H	H		3.51
I-8-54	3-CH <sub>3</sub>	5-Br	H	H		3.37
I-8-55	3-Cl	5-Br	H	H		2.94

### Preparation of starting materials of the formula (III)

#### Example (III-1)




(III-1)

5


A solution of 0.67 g (3 mmol) 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline, 0.80 g (3 mmol) of 2-bromo-3-chloro-5-(trifluoromethyl)pyridine and 0.05 g (0.68 mmol) of 1,1'-bis(diphenylphosphino)ferrocenepalladium(II)chloride in 15 ml dimethyl sulfoxide and 9 ml of a 2 M sodium carbonate solution were heated under an inert gas atmosphere for 16 h. For the work-up the reaction mixture was poured into water, extracted with ethyl acetate, dried over sodium sulphate and concentrated in vacuo. Purification via column chromatography yielded 0.79 g (2.9 mmol, 95 %) of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]aniline [log P (pH 2.3) = 2.79].

10

Example (III-2)

(III-2)

A mixture of 2.852 g (13 mmol) of 2-iodoaniline, 4.298 g of (17 mmol) of bis(pinacolato)diboron, 3.834 g (39 mmol) of potassium acetate and 24 mg (0.065 mmol) of bis(diphenylphosphino)ferrocenepalladium(II)chloride were stirred in 200 ml of dimethyl formamide at 80°C under argon. After 4 hours, 7 g (33 mmol) of 6-bromo-pyridine-3-carbaldehyde (*E*)-O-methyl-oxime (VII-1), 8.28 g (78 mmol) of sodium carbonate, 100 ml of water and another 24 mg (0.065 mmol) of bis(diphenylphosphino)ferrocenepalladium(II)chloride were added and the mixture was stirred for 12 hours at 80°C. The reaction mixture was cooled down to room temperature, poured into water, extracted with ethyl acetate, dried over sodium sulphate and concentrated under vacuum. Purification via column chromatography yielded 0.9 g (4.2 mmol, 32 %) of 6-(2-amino-phenyl)-pyridine-3-carbaldehyde (*E*)-O-methyl-oxime [ $\log P$  (pH 2.3) = 1.59].


15 Preparation of starting materials of the formula (IV)Example (IV-1)

(IV-1)

To a solution of 13.1 g (0.06 mol) o-iodoaniline and 12.1 g (0.12 mol) triethylamine in 250 ml tetrahydrofuran was added a solution of 15.0 g (0.07 mol) o-trifluoromethyl benzoic acid chloride in 250 ml tetrahydrofuran at 0°C. The reaction mixture was stirred for 30 min at 0°C and for 16 h at room temperature. Concentration in vacuo and column chromatography (ethyl acetate) yielded 23 g (0.06 mmol, 96 %) of *N*-(2-iodophenyl)-2-(trifluoromethyl)benzamide [ $\log P$  (pH 2.3) = 2.98].

**Preparation of starting materials of the formula (VII)****Example (VII-1)**

(VII-1)



5 10 g of 2-bromopyridine-5-carboxaldehyde (54 mmol) and 5.84 g (70 mmol) of *O*-methyl - hydroxylamine hydrochloride were dissolved in 100 ml of methanol and 50 ml of water. The mixture was stirred for 20 hours at room temperature. Methanol was then evaporated under reduced pressure, water added to the solid residue that was then filtered to afford 7 g (32 mmol) of 6-bromo-pyridine-3-carbaldehyde (*E*)-*O*-methyl-oxime [log P (pH 2.3) = 2.16].

10

The logP values given in the Preparation Examples were determined in accordance with EEC Directive 79/831 Annex V.A8 by HPLC (High Performance Liquid Chromatography) using a reversed-phase column (C 18). Temperature: 43°C.

15

Mobile phases for the determination in the acidic range: 0.1% aqueous phosphoric acid, acetonitrile; linear gradient from 10% acetonitrile to 90% acetonitrile.

20

Calibration was carried out using straight-chain alkan-2-ones (having 3 to 16 carbon atoms) with known logP values (determination of the logP values by the retention times using linear interpolation between two successive alkanones).

The lambda max values were determined in the maxima of the chromatographic signals using the UV spectra from 200 nm to 400 nm.

Use examplesExample A5 **Podosphaera test (apple) / protective**

Solvent:                    24.5    parts by weight of acetone  
                                 24.5    parts by weight of dimethylacetamide  
Emulsifier:                1      part by weight of alkylaryl polyglycol ether

10

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

15 To test for protective activity, young plants are sprayed with the preparation of active compound at the stated application rate. After the spray coating has dried on, the plants are inoculated with an aqueous spore suspension of the apple mildew pathogen *Podosphaera leucotricha*. The plants are then placed in a greenhouse at about 23°C and a relative atmospheric humidity of about 70 %.







20 Evaluation is carried out 10 days after the inoculation. 0 % means an efficacy which corresponds to that of the control, whereas an efficacy of 100 % means that no infection is observed.

Active compounds, application rates and test results are shown in the table below.









Table A


## Podosphaera test (apple) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
 (I-8-1)	100	100
 (I-8-3)	100	100
 (I-8-9)	100	97
 (I-6-7)	100	98
 (I-8-10)	100	100
 (I-8-11)	100	95

## Podospaera test (apple) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
 (I-8-12)	100	92
 (I-7-10)	100	100
 (I-8-27)	100	97
 (I-8-28)	100	96
 (I-8-20)	100	100
 (I-8-21)	100	100

## Podospaera test (apple) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
 (I-8-22)	100	100

Example B**Venturia Test (Apple) / protective**

- 5    Solvents:                    24.5    parts by weight acetone  
                                     24.5    parts by weight dimethylacetamide  
Emulsifier:                    1.0    part by weight alkylaryl polyglycol ether

10    To produce a suitable preparation of the active compound, 1 part by weight of active compound is mixed with the stated amounts of solvents and emulsifier, and the concentrate is diluted with water to the desired concentration.

15    To test for protective activity, young plants are sprayed with the preparation of active compound at the stated application rate. After the spray coating has dried on, the plants are inoculated with an aqueous conidia suspension of the apple scab pathogen *Venturia inaequalis* and then remain in an incubation cabin at about 20°C and 100 % relative atmospheric humidity for 1 day.







20    The plants are then placed in a greenhouse at about 21°C and a relative atmospheric humidity of about 90 %.

Evaluation is carried out 10 days after the inoculation. 0 % means an efficacy which corresponds to that of the control, whereas an efficacy of 100 % means that no infection is observed.








Active compounds, application rates and test results are shown in the table below.

Table B

## Venturia Test (Apple) / protective


Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
 (I-8-1)	100	100
 (I-8-3)	100	100
 (I-8-10)	100	100
 (I-8-11)	100	100
 (I-8-12)	100	100
 (I-8-14)	100	100

## Venturia Test (Apple) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
 (I-7-9)	100	100
 (I-7-10)	100	100
 (I-8-29)	100	98
 (I-8-27)	100	100
 (I-8-28)	100	98
 (I-8-22)	100	99
 (I-8-25)	100	99

**Venturia Test (Apple) / protective**

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
---	--	------------------



100

92

(I-8-30)

Example C**Alternaria test (tomato) / protective**

- 5    Solvent:                    49    parts by weight of N,N-dimethylformamide  
     Emulsifier:                1    part by weight of alkylaryl polyglycol ether

10    To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

15    To test for protective activity, young tomato plants are sprayed with the preparation of active compound at the stated application rate. 1 day after the treatment, the plants are inoculated with a spore suspension of *Alternaria solani* and then remain at 100 % rel. humidity and 20°C for 24 h. The plants then remain at 96 % rel. atmospheric humidity and a temperature of 20°C.






     Evaluation was carried out 7 days after the inoculation. 0 % means an efficacy which corresponds to that of the control, whereas an efficacy of 100 % means that no infection is observed.

20    Active compounds, application rates and test results are shown in the table below.



Table C

## Alternaria test (tomato) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
 (I-8-2)	750	100
 (I-8-3)	750	95
 (I-8-9)	750	95
 (I-7-6)	750	95
 (I-7-10)	750	100

Example D**Pyrenophora teres test (barley) / protective**

- 5    Solvent:                    50    parts by weight of N,N-dimethylacetamide  
    Emulsifier:                1    parts by weight of alkylaryl polyglycol ether

10    To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

15    To test for protective activity, young plants are sprayed with the preparation of active compound at the stated application rate. After the spray coating has dried on, the plants are sprayed with a conidia suspension of *Pyrenophora teres*. The plants remain in an incubation cabin at 20°C and 100 % relative atmospheric humidity for 48 hours.







    The plants are then placed in a greenhouse at a temperature of about 20°C and a relative atmospheric humidity of about 80 %.

20    Evaluation is carried out 8 days after the inoculation. 0 % means an efficacy which corresponds to that of the control, whereas an efficacy of 100 % means that no infection is observed.




    Active compounds, application rates and test results are shown in the table below.

Table D

## Pyrenophora teres test (barley) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
 (I-8-2)	500	100
 (I-8-3)	500	100
 (I-8-10)	500	100
 (I-7-4)	500	93
 (I-7-6)	500	93
 (I-8-13)	500	94

## Pyrenophora teres test (barley) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
 (I-8-14)	500	94
 (I-8-18)	500	94
 (I-7-5)	500	93

Example E**Puccinia test (wheat) / protective**

- 5 Solvent: 50 parts by weight of N,N-dimethylacetamide  
Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the  
10 desired concentration.

To test for protective activity, young plants are sprayed with the preparation of active compound at the stated application rate. After the spray coating has dried on, the plants are sprayed with a conidia suspension of *Puccinia recondita*. The plants remain in an incubation cabin at 20°C and 100 %  
15 relative atmospheric humidity for 48 hours.







The plants are then placed in a greenhouse at a temperature of approximately 20°C and a relative atmospheric humidity of 80 % to promote the development of rust pustules.

20 Evaluation is carried out 10 days after the inoculation. 0 % means an efficacy which corresponds to that of the control, whereas an efficacy of 100 % means that no infection is observed.

Active compounds, application rates and test results are shown in the table below.


Table E

Puccinia test (wheat) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
 (I-8-1)	500	100
 (I-8-2)	500	100
 (I-8-3)	500	100
 (I-8-10)	500	94
 (I-7-4)	500	100
 (I-7-6)	500	100

**Patent Claims**

## 1. Pyridinylanilides of the formula (I)



in which

R represents hydrogen, fluorine, chlorine, methyl or trifluoromethyl;

R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> independently of one another each represents hydrogen, halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

or represents in each case the grouping  $-C(Q^1)=N-Q^2$ , wherein

Q<sup>1</sup> represents hydrogen, hydroxyl or C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 9 identical or different halogen atoms, or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl and

Q<sup>2</sup> represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di(C<sub>1</sub>-C<sub>4</sub>-

alkyl)amino or phenyl; or represents C<sub>2</sub>-C<sub>4</sub>-alkenyloxy or C<sub>2</sub>-C<sub>4</sub>-alkynyloxy, represents phenyl, phenoxy, phenylthio, benzoyl, benzoylethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C<sub>1</sub>-C<sub>4</sub>-alkyl and C<sub>1</sub>-C<sub>4</sub>-alkoxy;

or

R<sup>2</sup> and R<sup>3</sup>, if attached to the pyridinyl moiety in ortho position to each other, furthermore together represent C<sub>3</sub>-C<sub>4</sub>-alkylene, C<sub>3</sub>-C<sub>4</sub>-alkenylene, C<sub>2</sub>-C<sub>3</sub>-oxyalkylene or C<sub>1</sub>-C<sub>2</sub>-dioxyalkylene, in each case optionally mono- to tetra-substituted, identically or differently, by fluorine, chlorine, oxo, methyl, ethyl, trifluoromethyl;

R<sup>4</sup> represents hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylsulfonyl, halogeno-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; (C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 7 fluorine-, chlorine- and/or bromine atoms, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl having in each case 1 to 6 fluorine-, chlorine- and/or bromine atoms, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl having in each case 1 to 13 fluorine-, chlorine- and/or bromine atoms; -COR<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,

R<sup>5</sup> represents hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>6</sub>-halogenoalkoxy, halogeno-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; or -COR<sup>10</sup>,

R<sup>6</sup> and R<sup>7</sup> independently of one another each represent hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>8</sub>-halogenoalkyl, halogeno-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halogenocycloalkyl having in each case 1 bis 9 fluorine-, chlorine- and/or bromine atoms,

R<sup>6</sup> and R<sup>7</sup> furthermore together with the nitrogen atom to which they are attached, represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and NR<sup>11</sup>, and which heterocycle may optionally be mono- to poly-substitu-



ted, identically or differently, by halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl,


R<sup>8</sup> and R<sup>9</sup> independently of one another each represent hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>8</sub>-halogenoalkyl, C<sub>3</sub>-C<sub>8</sub>-halogenocycloalkyl having in each case 1 bis 9 fluorine-, chlorine- and/or bromine atoms,

R<sup>8</sup> and R<sup>9</sup> furthermore together with the nitrogen atom to which they are attached, represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and NR<sup>11</sup>, and which heterocycle may optionally be mono- to poly-substituted, identically or differently, by halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl,

R<sup>10</sup> represents hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>6</sub>-halogenoalkoxy, halogeno-C<sub>1</sub>-C<sub>6</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms,

R<sup>11</sup> represents hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl,

A represents a radical of the formula (A1)



(A1), wherein


R<sup>12</sup> represents hydrogen, cyano, halogen, nitro, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy or C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio each having 1 to 5 halogen atoms, aminocarbonyl or aminocarbonyl-C<sub>1</sub>-C<sub>4</sub>-alkyl and

R<sup>13</sup> represents hydrogen, halogen, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-alkylthio and

R<sup>14</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, hydroxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkylthio-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl each having 1 to 5 halogen atoms, or phenyl,

or

A represents a radical of the formula (A2)




(A2), wherein

R<sup>15</sup> and R<sup>16</sup> independently of one another each represent hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub>-

alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms and  
 R<sup>17</sup> represents halogen, cyano or C<sub>1</sub>-C<sub>4</sub>-alkyl, or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl or C<sub>1</sub>-C<sub>4</sub>-  
 halogenoalkoxy each having 1 to 5 halogen atoms,

or

5 A represents a radical of the formula (A3)



R<sup>18</sup> and R<sup>19</sup> independently of one another each represent hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub>-  
 alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms and  
 R<sup>20</sup> represents hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1  
 10 to 5 halogen atoms,

or

A represents a radical of the formula (A4)



R<sup>21</sup> represents hydrogen, halogen, hydroxyl, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halo-  
 15 genoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy or C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio each having 1  
 to 5 halogen atoms,

or

A represents a radical of the formula (A5)



R<sup>22</sup> represents halogen, hydroxyl, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkyl-  
 20 thio, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio or C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy  
 each having 1 to 5 halogen atoms and

R<sup>23</sup> represents hydrogen, halogen, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkyl-  
 thio, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy each having 1 to 5 halogen  
 25 atoms, C<sub>1</sub>-C<sub>4</sub>-alkylsulphinyl or C<sub>1</sub>-C<sub>4</sub>-alkylsulphonyl,

or


A represents a radical of the formula (A6)



$R^{24}$  represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms and  
 $R^{25}$  represents  $C_1$ - $C_4$ -alkyl,  
 $Q^3$  represents a sulphur or oxygen atom, represents  $SO$ ,  $SO_2$  or  $CH_2$ ,  
 $p$  represents 0, 1 or 2, where  $R^{25}$  represents identical or different radicals if  $p$   
 represents 2,

or

A represents a radical of the formula (A7)




(A7), wherein

$R^{26}$  represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A8)




(A8), wherein

$R^{27}$  represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A9)




(A9), wherein

$R^{28}$  and  $R^{29}$  independently of one another each represent hydrogen, halogen, amino,  
 $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms and

$R^{30}$  represents hydrogen, halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1  
 to 5 halogen atoms,

or

A represents a radical of the formula (A10)




(A10), wherein

$R^{31}$  and  $R^{32}$  independently of one another each represent hydrogen, halogen, amino,  
 nitro,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms and

$R^{33}$  represents hydrogen, halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1  
 to 5 halogen atoms,

or

A represents a radical of the formula (A11)




(A11), wherein

$R^{34}$  represents hydrogen, halogen, amino,  $C_1$ - $C_4$ -alkylamino, di- $(C_1$ - $C_4$ -alkyl)-amino, cyano,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms and

$R^{35}$  represents halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A12)




(A12), wherein

$R^{36}$  represents hydrogen, halogen, amino,  $C_1$ - $C_4$ -alkylamino, di- $(C_1$ - $C_4$ -alkyl)-amino, cyano,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms and

$R^{37}$  represents halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A13)




(A13), wherein

$R^{38}$  represents halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A14)




(A14), wherein

$R^{39}$  represents hydrogen or  $C_1$ - $C_4$ -alkyl and

$R^{40}$  represents halogen or  $C_1$ - $C_4$ -alkyl,

or

A represents a radical of the formula (A15)




(A15), wherein

R<sup>41</sup> represents C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A16)




(A16), wherein

R<sup>42</sup> represents hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A17)



(A17), wherein

R<sup>43</sup> represents halogen, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio or C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy each having 1 to 5 halogen atoms,

excluded compounds of the formula (I), in which


R represents hydrogen and

R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> independently of one another each represents hydrogen, halogen; or straight-chain or branched alkyl having 1 to 4 carbon atoms; or straight-chain or branched halogenoalkyl having 1 to 4 carbon atoms; and

R<sup>4</sup> represents hydrogen

and

A represents a radical of the formula (A1)



(A1), wherein


R<sup>12</sup> represents halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl and

R<sup>13</sup> represents hydrogen and

R<sup>14</sup> represents methyl,

or

A represents a radical of the formula (A2)




(A2), wherein

$R^{15}$  and  $R^{16}$  independently of one another each represent hydrogen or  $C_1$ - $C_4$ -alkyl and  $R^{17}$  represents halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl,

5 or

A represents a radical of the formula (A4)




(A4), wherein

$R^{21}$  represents halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl,

or

10 A represents a radical of the formula (A5)




(A5), wherein

$R^{22}$  represents halogen and

$R^{23}$  represents hydrogen,

or

15 A represents a radical of the formula (A6)



(A6), wherein


$R^{24}$  represents methyl and

$Q^3$  represents a sulphur or  $CH_2$ ,

p represents 0,

20 or

A represents a radical of the formula (A9)




(A9), wherein

$R^{28}$  and  $R^{29}$  independently of one another each represent hydrogen or  $C_1$ - $C_4$ -alkyl and

$R^{30}$  represents methyl,

25 or

A represents a radical of the formula (A11)




(A11), wherein

$R^{34}$  represents hydrogen or  $C_1$ - $C_4$ -alkyl and

$R^{35}$  represents halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -halogenoalkyl,

or

5 A represents a radical of the formula (A16)



(A16), wherein

$R^{42}$  represents halogen.

2. Pyridinylanilides of the formula (I) according to Claim 1, in which

10 R represents hydrogen, fluorine, chlorine, methyl or trifluoromethyl;

$R^1$ ,  $R^2$  and  $R^3$  independently of one another each represents hydrogen, halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

15 or represents in each case straight-chain or branched alkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 4 carbon atoms and 1 to 9 identical or different halogen atoms;

20 or represents in each case straight-chain or branched alkylamino, dialkylamino, alkyl-carbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 4 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

25 or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms or  $C_3$ - $C_6$ -cycloalkyl and

$Q^2$  represents hydroxyl,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -halogenoalkyl or  $C_1$ - $C_4$ -halogenoalkoxy each having 1 to 9 identical or different halogen atoms,

30 or

$R^2$  and  $R^3$ , if attached to the pyridinyl moiety in ortho position to each other, furthermore together represent  $-(CH_2)_3-$ ,  $-(CH_2)_4-$ ,  $-CH=CH-CH=CH-$ ,  $-O(CH_2)_2-$ ,  $-O(CH_2)_3-$ ,

-OCH<sub>2</sub>O-, -O(CH<sub>2</sub>)<sub>2</sub>O-, in each case optionally mono- to tetra-substituted, identically or differently, by fluorine, chlorine, oxo, methyl, ethyl, trifluoromethyl,


- 5  $R^4$  represents hydrogen; C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylthio, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkylsulfonyl, halogeno-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; (C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 7 fluorine-, chlorine- and/or bromine atoms, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl having in each case 1 to 6 fluorine-, chlorine- and/or bromine atoms, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl having in each case 1 to 13 fluorine-, chlorine- and/or bromine atoms; -COR<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,
- 10  $R^5$  represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkoxy, halogeno-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; or -COR<sup>10</sup>,
- 15  $R^6$  and  $R^7$  independently of one another each represent hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, halogeno-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms,
- 20  $R^6$  and  $R^7$  furthermore together with the nitrogen atom to which they are attached, represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and NR<sup>11</sup>, and which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl,
- 25  $R^8$  and  $R^9$  independently of one another each represent hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl, C<sub>3</sub>-C<sub>6</sub>-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms,
- 30  $R^8$  and  $R^9$  furthermore together with the nitrogen atom to which they are attached, represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and NR<sup>11</sup>, and which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl,
- 35



$R^{10}$  represents hydrogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_3$ -alkoxy- $C_1$ - $C_3$ -alkyl,  $C_3$ - $C_6$ -cycloalkyl;  $C_1$ - $C_4$ -halogenoalkyl,  $C_1$ - $C_4$ -halogenoalkoxy, halogeno- $C_1$ - $C_3$ -alkoxy- $C_1$ - $C_3$ -alkyl,  $C_3$ - $C_6$ -halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms,

5  $R^{11}$  represents hydrogen or  $C_1$ - $C_4$ -alkyl,

A represents a radical of the formula (A1)



(A1), wherein

10  $R^{12}$  represents hydrogen, cyano, fluorine, chlorine, bromine, iodine, methyl, ethyl, iso-propyl, methoxy, ethoxy, methylthio, ethylthio, cyclopropyl,  $C_1$ - $C_2$ -halogenoalkyl,  $C_1$ - $C_2$ -halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms, trifluoromethylthio, difluoromethylthio, aminocarbonyl, aminocarbonylmethyl or aminocarbonylethyl and

15  $R^{13}$  represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, methoxy, ethoxy, methylthio or ethylthio and

$R^{14}$  represents hydrogen, methyl, ethyl, n-propyl, iso-propyl,  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms, hydroxymethyl, hydroxyethyl, cyclopropyl, cyclopentyl, cyclohexyl or phenyl,

or

20 A represents a radical of the formula (A2)




(A2), wherein

$R^{15}$  and  $R^{16}$  independently of one another each represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

25  $R^{17}$  represents fluorine, chlorine, bromine, cyano, methyl, ethyl,  $C_1$ - $C_2$ -halogenoalkyl or  $C_1$ - $C_2$ -halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

A represents a radical of the formula (A3)



R<sup>18</sup> and R<sup>19</sup> independently of one another each represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

5 R<sup>20</sup> represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

or


A represents a radical of the formula (A4)



10 R<sup>21</sup> represents hydrogen, fluorine, chlorine, bromine, iodine, hydroxyl, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl, C<sub>1</sub>-C<sub>2</sub>-halogenoalkoxy or C<sub>1</sub>-C<sub>2</sub>-halogenoalkylthio each having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

A represents a radical of the formula (A5)



15 R<sup>22</sup> represents fluorine, chlorine, bromine, iodine, hydroxyl, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio, C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms and

20 R<sup>23</sup> represents hydrogen, fluorine, chlorine, bromine, iodine, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, methoxy, ethoxy, methylthio, ethylthio, C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms, C<sub>1</sub>-C<sub>2</sub>-alkylsulphinyl or C<sub>1</sub>-C<sub>2</sub>-alkylsulphonyl,

or

25 A represents a radical of the formula (A6)




R<sup>24</sup> represents methyl, ethyl or C<sub>1</sub>-C<sub>2</sub>-halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

$R^{25}$  represents methyl or ethyl,  
 $Q^3$  represents a sulphur atom,  $SO_2$  or  $CH_2$ ,  
 $p$  represents 0 or 1,

or

5 A represents a radical of the formula (A9)



$R^{28}$  and  $R^{29}$  independently of one another each represent hydrogen, fluorine, chlorine, bromine, amino, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

10  $R^{30}$  represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

A represents a radical of the formula (A10)



15  $R^{31}$  and  $R^{32}$  independently of one another each preferably represent hydrogen, fluorine, chlorine, bromine, amino, nitro, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

$R^{33}$  represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

20 or

A represents a radical of the formula (A11)




25  $R^{34}$  represents hydrogen, fluorine, chlorine, bromine, amino,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino, cyano, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

$R^{35}$  represents fluorine, chlorine, bromine, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

A represents a radical of the formula (A12)




(A12), wherein

$R^{36}$  represents hydrogen, fluorine, chlorine, bromine, amino,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino, cyano, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

5  $R^{37}$  represents fluorine, chlorine, bromine, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

A represents a radical of the formula (A17)



(A17), wherein

10  $R^{43}$  preferably represents fluorine, chlorine, bromine, iodine, hydroxyl,  $C_1$ - $C_4$ -alkyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio,  $C_1$ - $C_2$ -halogenoalkyl or  $C_1$ - $C_2$ -halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms,


15 excluded compounds of the formula (I), in which

R represents hydrogen and

$R^1$ ,  $R^2$  and  $R^3$  independently of one another each represents hydrogen, halogen; or straight-chain or branched alkyl having 1 to 4 carbon atoms; or straight-chain or branched halogenoalkyl having 1 to 4 carbon atoms; and

20  $R^4$  represents hydrogen and

A represents a radical of the formula (A1)



(A1), wherein


25  $R^{12}$  represents fluorine, chlorine, bromine, iodine, methyl, ethyl, iso-propyl,  $C_1$ - $C_2$ -halogenoalkyl and

$R^{13}$  represents hydrogen and

$R^{14}$  represents methyl,

or

A represents a radical of the formula (A2)



(A2), wherein


$R^{15}$  and  $R^{16}$  independently of one another each represent hydrogen, methyl or ethyl  
and

$R^{17}$  represents fluorine, chlorine, bromine, methyl, ethyl, or  $C_1$ - $C_2$ -halogenoalkyl,

5

or

A represents a radical of the formula (A4)




(A4), wherein

$R^{21}$  represents fluorine, chlorine, bromine, iodine,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_2$ -halogenoalkyl,

10

or

A represents a radical of the formula (A5)



(A5), wherein


$R^{22}$  represents fluorine, chlorine, bromine, iodine and

$R^{23}$  represents hydrogen,

15

or

A represents a radical of the formula (A6)



(A6), wherein

$R^{24}$  represents methyl and


$Q^3$  represents a sulphur or  $CH_2$ ,

p represents 0,

20

or

A represents a radical of the formula (A9)



(A9), wherein


$R^{28}$  and  $R^{29}$  independently of one another each represent hydrogen, methyl or ethyl  
and

$R^{30}$  represents methyl,

25

or

A represents a radical of the formula (A11)



(A11), wherein

$R^{34}$  represents hydrogen, methyl or ethyl and

5  $R^{35}$  represents fluorine, chlorine, bromine, methyl, ethyl or  $C_1$ - $C_2$ -halogenoalkyl.

3. Pyridinylanilides of the formula (I) according to Claim 1, in which

R represents hydrogen, fluorine, chlorine, methyl or trifluoromethyl;

$R^1$ ,  $R^2$  and  $R^3$  independently of one another each represents hydrogen, fluorine, chlorine, bromine, cyano; methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxy, ethoxy, n- or iso-propoxy, n-, iso-, sec- or tert-butoxy, methylthio, ethylthio, n- or iso-propylthio, n-, iso-, sec- or tert-butylthio, trifluoromethyl, trifluoroethyl, difluoromethoxy, trifluoromethoxy, difluorochloromethoxy, trifluoroethoxy, cyclopropyl, cyclopentyl, cyclohexyl,

15 or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  represents hydrogen, methyl, ethyl, trifluoromethyl or cyclopropyl, and

$Q^2$  represents hydroxyl, methoxy, ethoxy, n-propoxy or iso-propoxy,

or

$R^2$  and  $R^3$ , if attached to the pyridinyl moiety in ortho position to each other, furthermore together represent  $-(CH_2)_3-$ ,  $-(CH_2)_4-$ ,  $-CH=CH-CH=CH-$ ,  $-OCH_2O-$ ,  $-O(CH_2)_2O-$ ,  $-OCF_2O-$ ,  $-O(CF_2)_2O-$ ,

$R^4$  represents hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, pentyl or hexyl, methylsulfinyl, ethylsulfinyl, n- or iso-propylsulfinyl, n-, iso-, sec- or tert-butylsulfinyl, methylsulfonyl, ethylsulfonyl, n- or iso-propylsulfonyl, n-, iso-, sec- or tert-butylsulfonyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl, trifluoromethyl, trichloromethyl, trifluoroethyl, difluoromethylthio, difluorochloromethylthio, trifluoromethylthio, trifluoromethylsulfinyl, trifluoromethylsulfonyl, trifluoromethoxymethyl;  $-CH_2-CHO$ ,  $-CH_2CH_2-CHO$ ,  $-CH_2CO-CH_3$ ,  $-CH_2CO-CH_2CH_3$ ,  $-CH_2CO-CH(CH_3)_2$ ,  $-CH_3CH_2CO-CH_3$ ,  $-CH_3CH_2CO-CH_2CH_3$ ,  $-CH_3CH_2CO-CH(CH_3)_2$ ,  $-CH_2-C(O)OCH_3$ ,  $-CH_2-C(O)OCH_2CH_3$ ,  $-CH_2-C(O)OCH(CH_3)_2$ ,  $-CH_3CH_2-C(O)OCH_3$ ,  $-CH_3CH_2-C(O)OCH_2CH_3$ ,  $-CH_3CH_2-C(O)OCH(CH_3)_2$ ,  $-CH_2CO-CF_3$ ,  $-CH_2CO-CCl_3$ ,  $-CH_2CO-CH_2CF_3$ ,  $-CH_2CO-CH_2CCl_3$ ,

30

$-\text{CH}_2\text{CH}_2-\text{CO}-\text{CH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CH}_2-\text{CO}-\text{CH}_2\text{CCl}_3$ ,  $-\text{CH}_2-\text{C}(\text{O})\text{OCH}_2\text{CF}_3$ ,  
 $-\text{CH}_2-\text{C}(\text{O})\text{OCF}_2\text{CF}_3$ ,  $-\text{CH}_2-\text{C}(\text{O})\text{OCH}_2\text{CCl}_3$ ,  $-\text{CH}_2-\text{C}(\text{O})\text{OCCl}_2\text{CCl}_3$ ,  
 $-\text{CH}_2\text{CH}_2-\text{C}(\text{O})\text{OCH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CH}_2-\text{C}(\text{O})\text{OCF}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CH}_2-\text{C}(\text{O})\text{OCH}_2\text{CCl}_3$ ,  
 $-\text{CH}_2\text{CH}_2-\text{C}(\text{O})\text{O}-\text{CCl}_2\text{CCl}_3$ ;  $-\text{COR}^5$ ,  $-\text{CONR}^6\text{R}^7$  or  $-\text{CH}_2\text{NR}^8\text{R}^9$ ,

5  $\text{R}^5$  represents hydrogen, methyl, ethyl, n- or iso-propyl, tert-butyl, methoxy, ethoxy, tert-butoxy, cyclopropyl; trifluoromethyl, trifluoromethoxy; or  $-\text{COR}^{10}$ ,

$\text{R}^6$  and  $\text{R}^7$  independently of one another each represent hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl; trifluoromethyl, trichloromethyl, trifluoroethyl, trifluoromethoxymethyl,

$\text{R}^6$  and  $\text{R}^7$  furthermore together with the nitrogen atom to which they are attached, represent a saturated heterocycle selected from the group consisting of morpholine, thiomorpholine and piperazine, which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by fluorine, chlorine, bromine or methyl and where the piperazine additionally at the second nitrogen atom may be substituted by  $\text{R}^{11}$ ,


$\text{R}^8$  and  $\text{R}^9$  independently of one another each represent hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl; trifluoromethyl, trichloromethyl, trifluoroethyl, trifluoromethoxymethyl,

$\text{R}^8$  and  $\text{R}^9$  furthermore together with the nitrogen atom to which they are attached, represent a saturated heterocycle selected from the group consisting of morpholine, thiomorpholine and piperazine, which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by fluorine, chlorine, bromine or methyl and where the piperazine additionally at the second nitrogen atom may be substituted by  $\text{R}^{11}$ ,

25  $\text{R}^{10}$  represents hydrogen, methyl, ethyl, n- or iso-propyl, tert-butyl, methoxy, ethoxy, n- or iso-propoxy, tert-butoxy, cyclopropyl; trifluoromethyl, trifluoromethoxy,

$\text{R}^{11}$  represents hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl,

A represents a radical of the formula (A1)



(A1), wherein

30

$\text{R}^{12}$  represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, iso-propyl, monofluoromethyl, monofluoroethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl, dichloromethyl, cyclopropyl,

methoxy, ethoxy, trifluoromethoxy, trichloromethoxy, methylthio, ethylthio, trifluoromethylthio or difluoromethylthio and


$R^{13}$  represents hydrogen, fluorine, chlorine, bromine, iodine or methyl and

$R^{14}$  represents hydrogen, methyl, ethyl, iso-propyl, trifluoromethyl, difluoromethyl, hydroxymethyl, hydroxyethyl or phenyl,

5

or

A represents a radical of the formula (A2)



(A2), wherein

$R^{15}$  and  $R^{16}$  independently of one another each represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl or trichloromethyl and


10

$R^{17}$  represents fluorine, chlorine, bromine, cyano, methyl, trifluoromethyl, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy or trichloromethoxy,

15

or

A represents a radical of the formula (A4)




(A4), wherein

$R^{21}$  represents hydrogen, fluorine, chlorine, bromine, iodine, hydroxyl, cyano, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy, trichloromethoxy, trifluoromethylthio, difluoromethylthio, difluorochloromethylthio or trichloromethylthio,

20

or

A represents a radical of the formula (A5)



(A5), wherein

$R^{22}$  represents fluorine, chlorine, bromine, iodine, hydroxyl, cyano, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio, tri-

30



fluoromethoxy, difluoromethoxy, difluorochloromethoxy or trichloromethoxy and

$R^{23}$  represents hydrogen, fluorine, chlorine, bromine, iodine, cyano, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl, methoxy, ethoxy, methylthio, ethylthio, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy, trichloromethoxy, methylsulphinyl or methylsulphonyl,

or

A represents a radical of the formula (A6)



$R^{24}$  represents methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl and


$R^{25}$  represents methyl,

$Q^3$  represents a sulphur atom or  $CH_2$ ,

$p$  represents 0,

or

A represents a radical of the formula (A9)



$R^{28}$  and  $R^{29}$  independently of one another each represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl and

$R^{30}$  represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl,

or

A represents a radical of the formula (A11)




$R^{34}$  represents hydrogen, fluorine, chlorine, bromine, amino, methylamino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl and

$R^{35}$  represents fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl,

or

A represents a radical of the formula (A17)



(A17), wherein

$R^{43}$  preferably represents fluorine, chlorine, bromine, iodine, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl,


excluded compounds of the formula (I), in which

R represents hydrogen and

$R^1$ ,  $R^2$  and  $R^3$  independently of one another each represents hydrogen, fluorine, chlorine, bromine; methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl; or trifluoromethyl or trifluoroethyl; and

$R^4$  represents hydrogen and

A represents a radical of the formula (A1)



(A1), wherein


$R^{12}$  represents fluorine, chlorine, bromine, iodine, methyl, ethyl, iso-propyl, monofluoromethyl, monofluoroethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl, dichloromethyl and

$R^{13}$  represents hydrogen and

$R^{14}$  represents methyl,

or

A represents a radical of the formula (A2)




(A2), wherein

$R^{15}$  and  $R^{16}$  independently of one another each represent hydrogen, methyl or ethyl and

$R^{17}$  represents fluorine, chlorine, bromine, methyl, ethyl, or trifluoromethyl,

or

A represents a radical of the formula (A4)




(A4), wherein

R<sup>21</sup> represents fluorine, chlorine, bromine, iodine, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl,

or

A represents a radical of the formula (A5)




(A5), wherein

R<sup>22</sup> represents fluorine, chlorine, bromine, iodine and

R<sup>23</sup> represents hydrogen,

or

A represents a radical of the formula (A6)



(A6), wherein


R<sup>24</sup> represents methyl and

Q<sup>3</sup> represents a sulphur or CH<sub>2</sub>,

p represents 0,

or

A represents a radical of the formula (A9)



(A9), wherein


R<sup>28</sup> and R<sup>29</sup> independently of one another each represent hydrogen, methyl or ethyl

and

R<sup>30</sup> represents methyl,

or

A represents a radical of the formula (A11)



(A11), wherein

$R^{34}$  represents hydrogen, methyl or ethyl and


$R^{35}$  represents fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

5 4. Pyridinylanilides of the formula (I) according to Claims 1, 2 or 3, in which  $R^4$  represents hydrogen.

5. Pyridinylanilides of the formula (I) according to Claims 1, 2 or 3, in which R represents hydrogen.

10

6. Pyridinylanilides of the formula (I-12)



in which

R,  $R^4$  and A are as defined in Claims 1, 2 or 3 and


15  $R^{1a}$  represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thio-carbamoyl;  
or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;  
20 or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;  
or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;  
25 or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;  
or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms  
30 in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;  
or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms, or  $C_3$ - $C_6$ -cycloalkyl and

$Q^2$  represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino or phenyl; or represents  $C_2$ - $C_4$ -alkenyloxy or  $C_2$ - $C_4$ -alkynyloxy, represents phenyl, phenoxy, phenylthio, benzoyl, benzoylethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched  $C_1$ - $C_4$ -alkyl and  $C_1$ - $C_4$ -alkoxy.

7. Pyridinylanilides of the formula (I-13)



(I-13)

in which

$R$ ,  $R^4$  and  $A$  are as defined in Claims 1, 2 or 3 and

$R^{1a}$  represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thio-carbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkyl-sulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;


or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms, or  $C_3$ - $C_6$ -cycloalkyl and

$Q^2$  represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino or phenyl; or represents  $C_2$ - $C_4$ -alkenyloxy or  $C_2$ - $C_4$ -alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoylethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched  $C_1$ - $C_4$ -alkyl and  $C_1$ - $C_4$ -alkoxy.

#### 8. Pyridinylanilides of the formula (I-14)



in which

$R$ ,  $R^4$  and  $A$  are as defined in Claims 1, 2 or 3 and

$R^{1a}$  and  $R^{2a}$  independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;


or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms, or  $C_3$ - $C_5$ -cycloalkyl and

$Q^2$  represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino or phenyl; or represents  $C_2$ - $C_4$ -alkenyloxy or  $C_2$ - $C_4$ -alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoyl-ethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched  $C_1$ - $C_4$ -alkyl and  $C_1$ - $C_4$ -alkoxy.

#### 9. Pyridinylanilides of the formula (I-15)




in which

R,  $R^4$  and A are as defined in Claims 1, 2 or 3 and

$R^{1a}$  and  $R^{2a}$  independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;  
 or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;  
 or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;  
 or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;  
 or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;  
 or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;  
 or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;  
 or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein  
 $Q^1$  represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms, or  $C_3$ - $C_6$ -cycloalkyl and  
 $Q^2$  represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino or phenyl; or represents  $C_2$ - $C_4$ -alkenyloxy or  $C_2$ - $C_4$ -alkynyloxy,  
 represents phenyl, phenoxy, phenylthio, benzoyl, benzoylphenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched  $C_1$ - $C_4$ -alkyl and  $C_1$ - $C_4$ -alkoxy.

#### 10. Pyridinylanilides of the formula (I-16)





in which

R, R<sup>4</sup> and A are as defined in Claims 1, 2 or 3 and

R<sup>1a</sup>, R<sup>2a</sup> and R<sup>3a</sup> independently of one another each represents halogen, cyano, nitro, amino,

hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

or represents the grouping -C(Q<sup>1</sup>)=N-Q<sup>2</sup>, wherein

Q<sup>1</sup> represents hydrogen, hydroxyl or C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 9 identical or different halogen atoms, or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl and

Q<sup>2</sup> represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino or phenyl; or represents C<sub>2</sub>-C<sub>4</sub>-alkenyloxy or C<sub>2</sub>-C<sub>4</sub>-alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoyl ethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclalkyl

having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C<sub>1</sub>-C<sub>4</sub>-alkyl and C<sub>1</sub>-C<sub>4</sub>-alkoxy.

- 5 11. Process for preparing pyridinylanilides of the formula (I) according to Claim 1, characterized in that


a) carboxylic acid derivatives of the formula (II)



in which

- 10 X¹ represents halogen or hydroxyl and  
A is as defined in Claim 1,

are reacted with amines of the formula (III)




15 in which

R, R¹, R², R³ and R⁴ are as defined in Claim 1,  
if appropriate in the presence of a catalyst, if appropriate in the presence of a condensing agent, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

20 or


b) halogeno-carboxamides of the formula (IV)



in which

- 25 R, R⁴ and A are as defined in Claim 1, and  
X² represents bromine or iodine,

are reacted with boronic acid derivatives of the formula (V)



in which


$R^1$ ,  $R^2$  and  $R^3$  are as defined in Claim 1, and

$A^1$  and  $A^2$  each represent hydrogen or together represent tetramethylethylene,

in the presence of a catalyst, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

c) carboxamide boronic acid derivatives of the formula (VI)




in which

$R$ ,  $R^4$  and  $A$  are as defined in Claim 1, and

$A^3$  and  $A^4$  each represent hydrogen or together represent tetramethylethylene,

are reacted with pyridinyl derivatives of the formula (VII)




in which

$R^1$ ,  $R^2$  and  $R^3$  are as defined in Claim 1,

in the presence of a catalyst, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

d) halogeno-carboxamides of the formula (IV)




in which

$R$ ,  $R^4$  and  $A$  are as defined in Claim 1, and

$X^2$  represents bromine or iodine,

are reacted with pyridinyl derivatives of the formula (VII)




in which

$R^1$ ,  $R^2$  and  $R^3$  are as defined in Claim 1,

in the presence of a palladium or platinum catalyst and in the presence of 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bis-1,3,2-dioxaborolane [bis(pinacolato)diboron], if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

e) pyridinylanilides of the formula (I-1)



in which

$R$ ,  $R^1$ ,  $R^2$ ,  $R^3$  and  $A$  are as defined in Claim 1,

are reacted with halogenides of the formula (VIII)




in which

$X^3$  represents chlorine, bromine or iodine,

$R^{4a}$  represents  $C_1$ - $C_8$ -alkyl,  $C_1$ - $C_6$ -alkylsulfinyl,  $C_1$ - $C_6$ -alkylsulfonyl,  $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -cycloalkyl;  $C_1$ - $C_6$ -halogenoalkyl,  $C_1$ - $C_4$ -halogenoalkylthio,  $C_1$ - $C_4$ -halogenoalkylsulfinyl,  $C_1$ - $C_4$ -halogenoalkylsulfonyl, halogeno- $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; formyl- $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -alkyl)carbonyl- $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -alkoxy)carbonyl- $C_1$ - $C_3$ -alkyl; ( $C_1$ - $C_3$ -halogenoalkyl)carbonyl- $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -halogenoalkoxy)carbonyl- $C_1$ - $C_3$ -alkyl having in each case 1 to 7 fluorine-, chlorine- and/or bromine atoms,


- (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl having in each case 1 to 6 fluorine-, chlorine- and/or bromine atoms, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl, (C<sub>1</sub>-C<sub>3</sub>-halogenoalkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-halogenoalkyl having in each case 1 to 13 fluorine-, chlorine- and/or bromine atoms; -COR<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,  
 5 R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined in Claim 1,  
 in the presence of a base and in the presence of a diluent.

12. Compositions for controlling unwanted microorganisms, characterized in that they comprise  
 10 at least one pyridinylanilide of the formula (I) according to Claim 1, in addition to extenders and/or surfactants.
13. Use of pyridinylanilides of the formula (I) according to Claim 1 for controlling unwanted  
 15 microorganisms.
14. Method for controlling unwanted microorganisms, characterized in that pyridinylanilides of the formula (I) according to Claim 1 are applied to the microorganisms and/or their habitats.
15. Process for preparing compositions for controlling unwanted microorganisms, characterized  
 20 in that pyridinylanilides of the formula (I) according to Claim 1 are mixed with extenders and/or surfactants.
16. Amines of the formula (III)



- 25 in which  
 R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are as defined in Claim 1,  
 excluded compounds of the formula (III), in which  
 R represents hydrogen and ^  
 R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> independently of one another each represents hydrogen, halogen, straight-chain  
 30 or branched alkyl having 1 to 4 carbon atoms or straight-chain or branched  
 halogenoalkyl having 1 to 4 carbon atoms; and R<sup>4</sup> represents hydrogen.

## 17. Amines of the formula




in which

R and R<sup>4</sup> are as defined in Claims 1, 2 or 3 and

- 5 R<sup>1a</sup> represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thio-carbamoyl;  
or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkyl-sulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;  
10 or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;  
or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;  
15 or represents in each case straight-chain or branched halogenoalkenyl or halogeno-alkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;  
or represents in each case straight-chain or branched alkylamino, dialkylamino, alkyl-carbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocar-bonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms  
20 in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;  
or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;  
or represents the grouping -C(Q<sup>1</sup>)=N-Q<sup>2</sup>, wherein  
25 Q<sup>1</sup> represents hydrogen, hydroxyl or C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1 to 9 identical or different halogen atoms, or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl and  
Q<sup>2</sup> represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino or phenyl; or represents C<sub>2</sub>-C<sub>4</sub>-alkenyloxy or C<sub>2</sub>-C<sub>4</sub>-alkynyloxy,  
30 represents phenyl, phenoxy, phenylthio, benzoyl, benzoylthenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclalkyl

having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C<sub>1</sub>-C<sub>4</sub>-alkyl and C<sub>1</sub>-C<sub>4</sub>-alkoxy.

5 18. Amines of the formula




in which

R and R<sup>4</sup> are as defined in Claims 1, 2 or 3 and

- 10 R<sup>1a</sup> represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thio-carbamoyl;  
or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkyl-sulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;  
or represents in each case straight-chain or branched alkenyl or alkenyloxy having in  
15 each case 2 to 6 carbon atoms;  
or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;  
or represents in each case straight-chain or branched halogenoalkenyl or halogeno-  
20 alkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;  
or represents in each case straight-chain or branched alkylamino, dialkylamino, alkyl-carbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocar-bonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms  
25 in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;  
or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;  
or represents the grouping -C(Q<sup>1</sup>)=N-Q<sup>2</sup>, wherein  
Q<sup>1</sup> represents hydrogen, hydroxyl or C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halogenoalkyl having 1  
30 to 9 identical or different halogen atoms, or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl and

Q<sup>2</sup> represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino or phenyl; or represents C<sub>2</sub>-C<sub>4</sub>-alkenyloxy or C<sub>2</sub>-C<sub>4</sub>-alkynyloxy, represents phenyl, phenoxy, phenylthio, benzoyl, benzoylphenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C<sub>1</sub>-C<sub>4</sub>-alkyl and C<sub>1</sub>-C<sub>4</sub>-alkoxy.

19. Amines of the formula



in which

R and R<sup>4</sup> are as defined in Claims 1, 2 or 3 and

R<sup>1a</sup> and R<sup>2a</sup> independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl; or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety; or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms; or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms; or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms; or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms



in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;


or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

5  $Q^1$  represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms, or  $C_3$ - $C_6$ -cycloalkyl and

$Q^2$  represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino or phenyl; or represents  $C_2$ - $C_4$ -alkenyloxy or  $C_2$ - $C_4$ -alkynyloxy,

10 represents phenyl, phenoxy, phenylthio, benzoyl, benzoylethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by  
15 halogen, in each case straight-chain or branched  $C_1$ - $C_4$ -alkyl and  $C_1$ - $C_4$ -alkoxy.

20. Amines of the formula



in which

20 R and  $R^4$  are as defined in Claims 1, 2 or 3 and

$R^{1a}$  and  $R^{2a}$  independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkyl-sulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

25 or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each  
30 case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;


or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms, or  $C_3$ - $C_6$ -cycloalkyl and

$Q^2$  represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino or phenyl; or represents  $C_2$ - $C_4$ -alkenyloxy or  $C_2$ - $C_4$ -alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoylthio, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched  $C_1$ - $C_4$ -alkyl and  $C_1$ - $C_4$ -alkoxy.

## 21. Amines of the formula



in which

R and R<sup>4</sup> are as defined in Claims 1, 2 or 3 and

R<sup>1a</sup>, R<sup>2a</sup> and R<sup>3a</sup> independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

or represents the grouping  $-C(Q^1)=N-Q^2$ , wherein

$Q^1$  represents hydrogen, hydroxyl or  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -halogenoalkyl having 1 to 9 identical or different halogen atoms, or  $C_3$ - $C_6$ -cycloalkyl and

$Q^2$  represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl)amino or phenyl; or represents  $C_2$ - $C_4$ -alkenyloxy or  $C_2$ - $C_4$ -alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoylthenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched  $C_1$ - $C_4$ -alkyl and  $C_1$ - $C_4$ -alkoxy.

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
20 January 2005 (20.01.2005)

PCT

(10) International Publication Number  
WO 2005/004606 A3

(51) International Patent Classification<sup>7</sup>: A01N 43/08,  
43/10, 43/40, 43/54, 43/56, 43/58, 43/60, 43/76, 43/78,  
C07D 213/61, 401/12, 417/12, 405/12, 411/12, 409/12

(74) Common Representative: BAYER CROPSCIENCE  
AG; Law and Patents, Patents and Licensing, 51368  
Leverkusen (DE).

(21) International Application Number:  
PCT/EP2004/007323

(22) International Filing Date: 5 July 2004 (05.07.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
03015733.3 10 July 2003 (10.07.2003) EP

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(71) Applicant (for all designated States except US): BAYER  
CROPSCIENCE AG [DE/DE]; Alfred-Nobel-Str. 50,  
40789 Monheim (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DUNKEL, Ralf  
[DE/DE]; Krischer Str. 22, 40789 Monheim (DE).  
ELBE, Hans-Ludwig [DE/DE]; Dasnöckel 59, 42329  
Wuppertal (DE). HARTMANN, Benoît [FR/DE]; Fär-  
berstr. 1, 40764 Langenfeld (DE). GREUL, Jörg, Nico  
[DE/DE]; Am Sandberg 30 a, 42799 Leichlingen (DE).  
WACHENDORFF-NEUMANN, Ulrike [DE/DE];  
Oberer Markenweg 85, 56566 Neuwied (DE). DAHMEN,  
Peter [DE/DE]; Aliebrückerstr. 61, 41470 Neuss (DE).  
KUCK, Karl-Heinz [DE/DE]; Pastor-Löh-Str. 30a,  
40764 Langenfeld (DE). MANSFIELD, Darren, James  
[GB/FR]; 25, Montee Bonafous, F-69004 Lyon (FR).  
COQUERON, Pierre-Yves [FR/FR]; 36, Rue de la Tete  
d'Or, F-69007 Lyon (FR). RIECK, Helko [DE/FR]; 9 rue  
Claude Monet, F-69110 Ste Foy les LYON (FR). DES-  
BORDES, Philippe [FR/FR]; 30, Rue Bancel, F-69004  
Lyon (FR).

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,  
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).


Published:

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

(88) Date of publication of the international search report:  
21 April 2005

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: PYRIDINYLANILIDES



(57) Abstract: Novel pyridinylanilides of the formula (I) in which R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and A are as defined in the description, a plurality of processes for preparing these substances and their use for controlling unwanted microorganisms, and also novel intermediates and their preparation.

# INTERNATIONAL SEARCH REPORT

International Application No.

PA/EP2004/0073 23

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>					
IPC 7	A01N43/08	A01N43/10	A01N43/40	A01N43/54	A01N43/56
	A01N43/58	A01N43/60	A01N43/76	A01N43/78	C07D213/61
	C07D401/12	C07D417/12	C07D405/12	C07D411/12	C07D409/12
According to International Patent Classification (IPC) or to both national classification and IPC					
<b>B. FIELDS SEARCHED</b>					
Minimum documentation searched (classification system followed by classification symbols)					
IPC 7 A01N C07D					
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched					
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)					
EPO-Internal, PAJ, WPI Data, EMBASE, BIOSIS, CHEM ABS Data, BEILSTEIN Data					
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>					
Category *	Citation of document, with indication, where appropriate, of the relevant passages				Relevant to claim No.
X	PATENT ABSTRACTS OF JAPAN. vol. 1996, no. 08, 30 August 1996 (1996-08-30) -& JP 08 092223 A (MITSUI TOATSU CHEM INC), 9 April 1996 (1996-04-09) cited in the application last paragraph in column 6 abstract; claims; table 1				1-16
Y	WO 01/53259 A (WALTER HARALD ; SYNGENTA PARTICIPATIONS AG (CH); SCHNEIDER HERMANN (DE) 26 July 2001 (2001-07-26) cited in the application page 97 - page 100; claim 2; examples 1.41-1.56, 1.100-1.117, 1.161-1.178, 1.233-1. 254; table 1				1-21
-/-					
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.					
* Special categories of cited documents : *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *C* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *Z* document member of the same patent family					
Date of the actual completion of the international search			Date of mailing of the international search report		
12 November 2004			28.02.2005		
Name and mailing address of the ISA European Patent Office, P.B. 5018 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016			Authorized officer  Härtinger, S		

Form: PCT/ISA/210 (second sheet) (January 2004)

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP2004/007323

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95/09846 A (DU PONT ; DENES LUCIAN RADU (US)) 13 April 1995 (1995-04-13) cited in the application	1-16
Y	page 30, line 23; example 48; table 1 page 17, line 22; claims 4,6-10	17-21
Y	WO 96/16954 A (AGREVO UK LTD ; RIORDAN PETER DOMINIC (GB); WEST PETER JOHN (GB); BODD) 6 June 1996 (1996-06-06) claim 1; examples 22,23,26,27,33,35,42	1-21
Y	EP 0 545 099 A (BASF AG) 9 June 1993 (1993-06-09) cited in the application page 4, line 1 - line 16; claim 1 page 2, line 1 - line 27 page 52	1-21
Y	WO 93/11117 A (MONSANTO CO) 10 June 1993 (1993-06-10) cited in the application claim 1	1-16

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP2004/007323

## Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  

Although claims 13 and 14 have been drafted such that they may also embrace a method of treatment of the human or animal body, the search has been carried out and based on the alleged anti-microbiological effects of the compound.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-15 (part)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ASA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-15

A is a 5-membered heteroaromatic ring with 2 heteroatoms in 1,2 position  
---

2. claims: 1-15

A is a 5-membered heteroaromatic ring with 2 heteroatoms in 1,3 position  
---

3. claims: 1-15

A is a 5-membered heteroaromatic ring with 1 heteroatom  
---

4. claims: 1-15

A is a benzene ring  
---

5. claims: 1-15

A is a 6-membered heteroaromatic ring with 1 heteroatom  
---

6. claims: 1-15

A is a 6-membered heteroaromatic ring with 2 heteroatoms  
---

7. claims: 1-15

A is a 6-membered partially unsaturated heterocyclic ring  
---

8. claims: 1-15

A is a 5-membered partially unsaturated heterocyclic ring  
---

9. claims: 16-21

intermediates of the general formula III  
---



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/007323

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 08092223	A	09-04-1996	NONE	
WO 0153259	A	26-07-2001	AU 772635 B2	06-05-2004
			AU 3543301 A	31-07-2001
			BR 0107738 A	22-10-2002
			CA 2397008 A1	26-07-2001
			CN 1427823 T	02-07-2003
			CZ 20022496 A3	16-10-2002
			WO 0153259 A1	26-07-2001
			EP 1252140 A1	30-10-2002
			HU 0203960 A2	28-03-2003
			JP 2003520269 T	02-07-2003
			PL 357312 A1	26-07-2004
			US 2004049035 A1	11-03-2004
			US 2004106521 A1	03-06-2004
			ZA 200205641 A	03-11-2003
WO 9509846	A	13-04-1995	AU 7834494 A	01-05-1995
			CA 2173326 A1	13-04-1995
			EP 0722441 A1	24-07-1996
			WO 9509846 A1	13-04-1995
			US 5631206 A	20-05-1997
WO 9616954	A	06-06-1996	AU 4302896 A	19-06-1996
			WO 9616954 A1	06-06-1996
			EP 0794950 A1	17-09-1997
			ZA 9510223 A	29-07-1996
EP 0545099	A	09-06-1993	AT 149487 T	15-03-1997
			AU 656243 B2	27-01-1995
			AU 2855492 A	27-05-1993
			CA 2081935 A1	23-05-1993
			CZ 9203448 A3	13-10-1993
			CZ 289478 B6	16-01-2002
			DE 59208113 D1	10-04-1997
			DK 545099 T3	24-03-1997
			EP 0545099 A2	09-06-1993
			ES 2098421 T3	01-05-1997
			GR 3023336 T3	29-08-1997
			HU 62861 A2	28-06-1993
			IL 103614 A	24-09-1998
			JP 3202079 B2	27-08-2001
			JP 5221994 A	31-08-1993
			JP 2001253802 A	18-09-2001
			JP 2001316210 A	13-11-2001
			KR 267518 B1	16-10-2000
			NZ 245194 A	27-02-1996
			PL 296677 A1	18-10-1993
			SK 344892 A3	08-03-1995
			US 5480897 A	02-01-1996
			US 5556988 A	17-09-1996
			US 5589493 A	31-12-1996
			US 5330995 A	19-07-1994
			ZA 9208977 A	19-05-1994
WO 9311117	A	10-06-1993	US 5223526 A	29-06-1993
			AR 248397 A1	18-08-1995
			AT 149490 T	15-03-1997

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PSA/EP2004/007323

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9311117 A		AU 657598 B2	16-03-1995
		AU 3240793 A	28-06-1993
		BR 9206869 A	28-11-1995
		CA 2123122 A1	10-06-1993
		CZ 9401260 A3	15-02-1995
		DE 69217997 D1	10-04-1997
		DE 69217997 T2	04-09-1997
		EP 0623113 A1	09-11-1994
		HU 67795 A2	28-04-1995
		JP 7501549 T	16-02-1995
		LT 234 A	15-06-1994
		MD 940085 A	30-06-1995
		MX 9207038 A1	01-06-1993
		NZ 246269 A	26-07-1995
		SI 9200352 A	30-06-1993
		SK 67894 A3	08-02-1995
		TR 26592 A	15-03-1995
		WO 9311117 A1	10-06-1993
		ZA 9209441 A	25-08-1993
		ZW 18792 A1	12-05-1993
		CN 1078234 A	10-11-1993